


# Xeriscape Conversion Study



SOUTHERN NEVADA  
WATER AUTHORITY

## Final Report

2005

By

Kent A. Sovocool  
Senior Conservation Programs Analyst  
Southern Nevada Water Authority

With data processing assistance from

Mitchell Morgan  
Assistant Management Analyst  
Southern Nevada Water Authority

Funded in part by a grant from  
Bureau of Reclamation  
U.S. Department of Interior

### SNWA Member Agencies

- *Big Bend Water*
- *City of Boulder City*
- *City of Las Vegas*
- *City of Henderson*
- *City of North Las Vegas*
- *Clark County Water Reclamation District*
- *Las Vegas Valley Water District*

## Table of Contents

TABLE OF TABLES.....	ii
TABLE OF FIGURES.....	iii
ABSTRACT.....	4
ACKNOWLEDGEMENTS.....	6
INTRODUCTION AND BACKGROUND.....	7
<i>XERISCAPE AND WHAT IT MAY MEAN FOR WATER CONSERVATION</i> .....	7
<i>NEVADA’S COLORADO RIVER RESOURCES AND THE SPECIAL IMPORTANCE OF OUTDOOR WATER</i> .....	8
<i>CONSERVATION</i> .....	8
<i>THE RESEARCH STUDY</i> .....	10
METHODOLOGY.....	11
<i>STUDY GROUPS AND MONITORING</i> .....	11
<i>GENERAL DATA METHODS, STRATEGIES, AND STATISTICS</i> .....	13
<i>PRE/POST ANALYSES</i> .....	13
<i>ANALYSES OF SAVINGS OVER TIME AND SEASONS</i> .....	14
<i>COMPARATIVE PER-UNIT AREA IRRIGATION ANALYSES</i> .....	18
<i>MULTIVARIATE ANALYSES TO IDENTIFY SIGNIFICANT SOURCES OF VARIABILITY</i> .....	20
<i>ECONOMIC ANALYSES</i> .....	21
RESULTS AND DISCUSSION.....	22
<i>REDUCTION IN TOTAL HOUSEHOLD WATER CONSUMPTION FOLLOWING CONVERSION</i> .....	22
<i>TO XERISCAPE</i> .....	22
<i>ASSESSMENT OF SAVINGS POTENTIAL ACROSS TIME AND SEASONS</i> .....	24
<i>COMPARISON OF PER UNIT AREA WATER APPLICATION BETWEEN TURFGRASS AND</i> .....	30
<i>XERIC LANDSCAPE</i> .....	30
Annual application.....	30
Monthly Application.....	32
<i>SOURCES OF SIGNIFICANT VARIABILITY IN SINGLE-FAMILY RESIDENTIAL</i> .....	40
<i>CONSUMPTION</i> .....	40
Variability in Annual Residential Consumption.....	41
Variability in Annual Consumption for Irrigation of Monitored Xeric Landscape.....	46
<i>FINANCIAL SAVINGS ASSOCIATED WITH CONVERSION PROJECTS AND COST EFFICIENCY</i> .....	50
<i>ESTIMATED APPROPRIATE LEVEL OF FINANCIAL INCENTIVE</i> .....	54
Homeowner Perspective.....	54
SNWA Perspective.....	58
EXECUTIVE SUMMARY AND CONCLUSIONS.....	60
REFERENCES.....	64
APPENDICES.....	66
<i>Appendix 1: Attachment A (Scope of Work) for BOR</i> .....	67
<i>Agreement 5-FC-30-00440 as Revised 11/19/98</i> .....	67
<i>Appendix 2: Multivariate Model Details</i> .....	73

<i>Appendix 3: Raw Data</i> .....	74
<i>Appendix 4: Information on Single-Family Residential Water Bill Model</i> .....	93
<i>Appendix 5: Information on Homeowner Perspective Model</i> .....	94
<i>Appendix 6: Information on SNWA’s Water Smart Landscapes Program</i> .....	96

## **Table of Tables**

TABLE 1: Planned Pre-/Post-Retrofit Analyses for XS Group.....	14
TABLE 2: Planned Pre-/Post-Retrofit Analyses for TS Group.....	14
TABLE 3: Planned Post-Retrofit Analyses for XS Group Across Time.....	15
TABLE 4: Planned Post-Retrofit Analyses for TS Group Across Time.....	15
TABLE 5: Enhanced Post-Retrofit Analyses for XS Group Across Time.....	16
TABLE 6: Enhanced Post-Retrofit Analyses for TS Group Across Time.....	16
TABLE 7: Planned Summer Post-Retrofit Analyses for XS Group.....	17
TABLE 8: Planned Summer Post-Retrofit Analyses for TS Group.....	18
TABLE 9: Planned Comparative Analysis of Turf and Xeric Per Unit.....	19
TABLE 10: Planned Comparative Analysis of Turf and Xeric Per Unit.....	19
TABLE 11: Pre-/Post-Retrofit Analyses for XS Group.....	22
TABLE 12: Pre-/Post-Retrofit Analyses for TS Group.....	23
TABLE 13: Enhanced Post-Retrofit Analyses for XS Group Across Time.....	26
TABLE 14: Summer Post-Retrofit Analyses for XS Group.....	27
TABLE 15: Enhanced Post-Retrofit Analyses for TS Group Across Time.....	28
TABLE 16: Summer Post-Retrofit Analyses for TS Group.....	29
TABLE 17: Annual Per-Unit Area Application to Turf and Xeriscape.....	30
TABLE 18: Monthly Per-Unit Area Application to Turf and Xeriscape.....	33
TABLE 19: Multivariate Regression Model of Annual Single-Family Residential.....	73
TABLE 20: Multivariate Regression Model of Annual Xeric Study Area Consumption.....	73

## Table of Figures

FIGURE 1: Pre-/Post-Retrofit Consumption for XS and Comparison Groups .....	22
FIGURE 2: Pre-/Post-Retrofit Consumption for XS Group Across Time .....	25
FIGURE 3: Annual Per Unit Area Application to Turf and Xeriscape .....	31
FIGURE 4: Distribution of Annual Per Unit Area Application Data for Turf and Xeriscape .....	32
FIGURE 5: Monthly Per-Unit Area Application for Turf and Xeric Areas .....	34
FIGURE 6: Monthly Per-Unit Area Savings (Turf Area Application– Xeric Area Application) .....	35
FIGURE 7: Monthly Per-Unit Area Application to Turf and Reference.....	36
FIGURE 8: Monthly Per-Unit Area Application to Xeric Areas .....	36
FIGURE 9: Absolute Departure in Irrigation Application from Derived Respective .....	37
FIGURE 10: Relative Departure in Irrigation Application from Derived Respective .....	38
FIGURE 11: Relative Departure in Irrigation Application from Derived Respective .....	39
FIGURE 12: Average Monthly Maintenance Time and Annual Direct Expenditures.....	51
FIGURE 13: Modeled Monthly Water Bill for a Typical Las Vegas Area Home and .....	52
FIGURE 14: Modeled Monthly Water Bill Savings for A Typical Las Vegas Area .....	53
FIGURE 15: Summary of Economics of Typical Single-Family Xeriscape.....	57

## Abstract

The authors present a manuscript covering the Southern Nevada Water Authority's (SNWA) multi-year Xeriscape Conversion Study, which was funded in part by the Bureau of Reclamation - Lower Colorado Regional Area<sup>1</sup>.

Xeriscape (low-water-use landscaping) has held the promise of significant water savings for a number of years, but how much exactly it can save, especially as a practical residential landscape concept has been a point of debate and conjecture. Lacking to date has been a truly experimental quantitative study in which per-unit area application data has been gathered to quantify savings estimates (for a variety of reasons, most research has been limited to the total household level, with comparisons involving homes that are mostly xeriscape or traditional landscaping). Recognizing the need for more exacting (and locally applicable) savings estimates, SNWA conducted a study that could yield quantitative savings estimates of what a xeriscape conversion facilitation program could yield under real world conditions.

The experimental study involved recruiting hundreds of participants into treatment groups (a Xeric Study and a Turf Study Group and control groups), as well as the installation of submeters to collect per unit area application data. Data on both household consumption and consumption through the submeters was collected, as well as a wealth of other data. In most cases, people in the xeric study group converted from turf to xeriscape, though in some cases recruitment for this group was enhanced by permitting new landscapes with xeric areas suitable for study to be monitored. Portions of xeric areas were then submetered to determine per-unit area water application for xeric landscapes. The TS Group was composed of more traditional turfgrass-dominated landscapes, and submeters were installed to determine per-unit area application to these areas as well. Submeter installation, data collection, and analysis for a small side-study of multi-family/commercial properties also took place.

Results show a significant average savings of 30% (96,000 gallons) in total annual residential consumption for those who converted from turf to xeriscape. The per-unit area savings as revealed by the submeter data was found to be 55.8 gallons per square foot (89.6 inches precipitation equivalents) each year. Results showed that savings yielded by xeriscapes were most pronounced in summer. A host of other analyses covering everything from the stability of the savings to important factors influencing consumption, to cost effectiveness of a xeriscape conversion program are contained within the report.

**An abbreviated summary of the report's findings appears as the *Executive Summary and Conclusions* section (pg. 60).**

---

<sup>1</sup>This report with written and electronic appendices satisfies a deliverables requirement pursuant the applicable funding agreement with the Bureau of Reclamation (Cooperative Agreement #5-FC-30-00440). SNWA gratefully acknowledges BOR for its funding assistance with this project.

Keywords: *water conservation, xeriscape, xeric, landscape conversion, desert landscape low-water-use, plants, landscape, irrigation, residential water consumption, outdoor water use, submeter, irrigation controller, resource conservation, incentive programs, utility, water resources.*

## Acknowledgements

There have over the course of this study been so many contributors to this research that thanking each of them individually is an impossibility; however, the authors would like to express their gratitude to individuals in the following groups and organizations without whom this research would not have been possible.

- The study participants without whom no data could have been collected.
- The Southern Nevada Water Authority (SNWA) and the numerous personnel who have supported and promoted this research. The authors especially thank members of the Conservation Division, present and past, which have supported this research over the years.
- The member agencies of the SNWA listed on the cover. Special thanks go to Boulder City, the cities of Henderson and North Las Vegas, and the Las Vegas Valley Water District for the reading of meters and submeters related to the study.
- The SNWA Xeriscape Conversion Study Team and all those who have contributed to study recruitment, data collection, and data processing.
- Aquacraft, Inc. for preliminary analyses and collection of trace flow data.
- The Bureau of Reclamation Lower Colorado River Office and members of this office who helped facilitate funding for the research.
- Those whom have reviewed and commented on the Xeriscape Conversion Study and the results. The authors graciously acknowledge the contributions of reviewers of the final manuscript. Final manuscript reviewers included SNWA personnel in the Conservation and Resources Divisions, members of the Las Vegas Valley Water District's Resources Division, a contractor to the SNWA, and outside reviewers at the City of Austin, Texas. The Bureau of Reclamation also had the opportunity to comment on a draft prelude to final submission.
- Those organizations that have acted as a forum for exhibiting this research.
- Those who have worked for the study in the past and have gone on to other things.
- Those who have supported the research and the research personnel in other capacities.

# Introduction and Background

## **XERISCAPE AND WHAT IT MAY MEAN FOR WATER CONSERVATION**

In the Mojave Desert of the southwestern United States, typically 60 to 90% of potable water drawn by single-family residences in municipalities is used for outdoor irrigation. Thus, in this region, and indeed most of the entire Southwest, the most effective conservation measures are oriented towards reducing outdoor water consumption. A commonly considered method for accomplishing water conservation is to use xeriscape (low-water-use landscaping) in place of traditional turf. Xeriscape is based on seven principles:

- Sound Landscape Planning and Design
- Limitation of Turf to Appropriate Areas
- Use of Water-efficient Plants
- Efficient Irrigation
- Soil Amendments
- Use of Mulches
- Appropriate Landscape Maintenance

The term “xeriscape” was invented by Nancy Leavitt, of Denver Water (a public utility) in the early 1980s as a fusion of the Greek word “xeros” (meaning dry or arid) and landscape. Denver Water trademarked the term shortly thereafter though it has entered the English vernacular over the last 20 years as the concept has spread globally.

So promising was xeriscape, that water purveyors and others interested in conservation began actively promoting xeriscape in place of traditional landscape as early as the mid-80s as part of water conservation strategies. The need to better understand its true effectiveness as a conservation tool led to a host of studies being conducted in the 1990s, which have generally pegged savings associated with xeriscape at between 25% and 42% for the residential sector (Bent<sup>1</sup> 1992, Testa and Newton<sup>2</sup> 1993, Nelson<sup>3</sup> 1994, Gregg<sup>4</sup> et al. 1994). The variation in savings estimates is due to a large number of factors ranging from the different climates of each study locality, different local definitions of xeriscape, and different study methodologies employed.

The work done to this point has greatly advanced the water conservation community’s ability to evaluate, modify, and justify programs to encourage the use of xeriscaping as an integral component of water conservation plans. Utilities, water districts, cities, counties, and states are beginning to promote xeriscape as a cost-effective, mutually beneficial alternative to traditional turfgrass-dominated landscapes. Recently, this interest has increased at the national level, and this study is part of that evolution. Interest is further enhanced at the time of publication of this report due to a significant drought impacting the Colorado River Basin and much of the western United States.



## **NEVADA’S COLORADO RIVER RESOURCES AND THE SPECIAL IMPORTANCE OF OUTDOOR WATER CONSERVATION**

The Colorado River serves as the lifeblood for many of the communities of the southwestern United States, permitting society to flourish, despite the harsh, arid conditions that often define it. It serves the needs of millions within the region and its yearly volume is entirely divided up by the Colorado River Compact<sup>5</sup> and subsequent legislation and legal decisions, known as the “Law of the River” that specify allocations for each of the states (and Mexico) through which it flows. Among other things, the Bureau of Reclamation – Lower Colorado Region (BOR-LCR) is charged with maintaining an adequate and established allocation of water for each of the states in the arid Lower Basin. Since water demand management is ultimately accomplished at local levels, BOR-LCR actively partners with entities that divert Colorado River water to encourage conservation. In southern Nevada, the major regional organization meeting this criterion is the Southern Nevada Water Authority (SNWA).

In 1991 the SNWA was established to address water on a cooperative local basis, rather than an individual water purveyor basis. The SNWA is committed to managing the region’s water resources and developing solutions that ensure adequate future water supplies for southern Nevada. The member agencies are the cities of Boulder City, Henderson, Las Vegas, North Las Vegas, the Big Bend Water District, the Clark County Water Reclamation District, and the Las Vegas Valley Water District. As southern Nevada has grown into a metropolitan area and a world-famous vacation destination, so too have its water needs. The SNWA was created to plan and provide for the present and future water needs of the area.


Five different water purveyors provide potable water to most of Clark County. Big Bend Water District provides water to the community of Laughlin; the cities of Boulder City and Henderson provide water to their respective communities. The Las Vegas Valley Water District provides water to the City of Las Vegas and portions of unincorporated Clark County; the City of North Las Vegas provides water within its boundaries and to adjacent portions of unincorporated Clark County and the City of Las Vegas. The SNWA member agencies serve approximately 96% of the County’s population.

Southern Nevada’s climate is harsh. The Las Vegas Valley receives only 4.5 inches of precipitation annually on average, has a yearly evapotranspirational (ET) water requirement of nearly 90 inches, and it is one of the fastest growing metropolitan areas in the United States. Clark County, the southernmost county in Nevada, has a population in excess of 1.6 million people and has been experiencing extremely strong economic growth in recent years with correspondent annual population growth averaging in excess of 5% percent. The primary economic driver of Clark County’s economy is the tourism and gaming industry, with an annual visitor volume in excess of 30 million people per year. Today more than 7 out of every 10 Nevadans call Clark County home.

Consumptive use (use where Colorado River water does not return to the Colorado River) is of paramount interest to SNWA (specifically, consumptive use is defined by SNWA as

the summation of yearly diversions minus the sum of return flows to the River). A 1964 Supreme Court Decree in *Arizona v. California* verified the Lower Basin apportionment of 7.5 million acre feet (MAF) among Arizona, California, and Nevada, including Nevada's consumptive use apportionment of 300,000 acre feet per year (AFY) of Colorado River water as specified initially in the Colorado River Compact<sup>5</sup> and Boulder Canyon Project Act<sup>6</sup>. Return flows in Nevada consist mainly of highly treated Colorado River wastewater that is returned to Lake Mead and to the Colorado River at Laughlin, Nevada. With return flow credits, Nevada can actually divert more than 300,000 AFY, as long as the consumptive use is no more than 300,000 AFY (see diagram below). Since Colorado River water makes up roughly 90% of SNWA's current water-delivering resource portfolio, it means that in terms of demand management, reduction of water used outdoors (i.e., water unavailable for accounting as return flow) is much more important in terms of extending water resources than reduction of indoor consumption at this point in time.

**Diagram Showing Dynamic of Diversions, Return Flow Credits (from indoor uses) and Consumptive Use**



Since most of the SNWA (Authority) service area contains relatively scarce local reserves (there are little surface or groundwater resources) and since, as explained above, its Colorado River apportionment is limited, the organization has an aggressive conservation program that began in the 1990s. The Authority has been committed to achieving a 25% level of conservation (versus what consumption would have been without conservation) by the year 2010 (note though that soon this goal will be revised to probably be even more aggressive in the immediate future due to the drought). In 1995, the SNWA member agencies entered into a Memorandum of Understanding (MOU) regarding a regional water conservation plan. The MOU, updated in 1999, identifies specific management practices, timeline, and criteria the member agencies agree to follow in order to implement water conservation and efficiency measures.

The programs or Best Management Practices (BMPs) listed in the MOU include water measurement and accounting systems; incentive pricing and billing; water conservation coordinators; information and education programs; distribution system audit programs; customer audit and incentive programs; commercial and industrial audit and incentive programs; landscape audit programs; landscape ordinances; landscape retrofit incentive programs; waste-water management and recycling programs; fixture replacement programs; plumbing regulations, and water shortage contingency plans. The BMPs provide the framework for implementing the water conservation plan and guidance as to the methods to be employed to achieve the desired savings.

## **THE RESEARCH STUDY**

The potentially large water savings attainable with the broad-scale use of xeriscaping and the fact that associated reductions are in consumptive-use water makes xeriscape of paramount interest for both BOR and SNWA. For this reason, a partnership between BOR and SNWA was formed to investigate the savings that could be obtained with a program to encourage converting traditional turfgrass landscape to xeriscape. This was formally implemented as a Cooperative Agreement<sup>7</sup> in 1995. With its signing, a multi-year study of xeriscape was born, which has come to be known as the SNWA Xeriscape Conversion Study (XCS). As delineated in the most recent version of the Scope (Appendix 1) for this agreement, the objectives of the Study are to:

- Objective 1: Identify candidates for participation in the Study and monitor their water use.
- Objective 2: Measure the average reduction in water use among Study participants.
- Objective 3: Measure the variability of water savings over time and across seasons.
- Objective 4: Assess the variability of water use among participants and to identify what factors contribute to that variability.
- Objective 5: Measure the capital costs and maintenance costs of landscaping among participants.
- Objective 6: Estimate incentive levels necessary to induce a desired change in landscaping.

SNWA assembled a team to support the XCS, and field data was collected through 2001 with the draft final report finished in 2004 (intermediate reports outlined some of the major conclusions). By agreement, the SNWA agreed to provide the raw data collected for possible use in national research efforts by BOR (data was included with the final version of this manuscript submitted to BOR).

# Methodology

## STUDY GROUPS AND MONITORING

The study team recruited participants who live in single-family residences within the following entities' water jurisdictions: The Las Vegas Valley Water District (77% of the participants in the entire study group), Henderson (12%), North Las Vegas (9%), and Boulder City (2%).

There are a total of three groups in the XCS, the Xeriscape Study (XS) Group, the Turf Study (TS) Group, and a non-contacted Comparison Group. The XS Group is composed of residents who converted at least 500 square feet (sqft) of traditional turfgrass to xeric landscape as well as residents who installed new xeric landscaping. To clarify, in this region, xeric landscaping is principally composed of a combination of desert-adapted shrubs, trees, some ornamental grasses, and mulch (often rock). A \$0.45 per square foot incentive helped the property owner by absorbing some, but not the majority, of the cost of the conversion. Homeowners were required to plant sufficient vegetation so that the xeric landscape would at a minimum have 50% canopy coverage at maturity. This avoided the creation of unattractive "zeroscapes" composed exclusively of rocks, which could potentially act as urban heat islands. The incentive was capped for each residence at \$900 for 2,000 sqft; however, many residents converted more landscape than that which qualified for the incentive with the cap. Indeed, the average area converted in this study group was 2,162 sqft. A total of 472 properties were enrolled in the Study as XS Group participants. Aerial photographs, supported by ground measures, were used for recording areas. As a supplement to the main experimental group, 26 multi-family and commercial properties were submetered as well.

In return for the incentive, XS Group residents agreed to ongoing monitoring of their water consumption. This was accomplished in two ways. First, mainmeter data was taken from standard monthly meter reading activity (this was for assessing water use at the entire single-family residence level). Second, residents agreed to installation of a submeter that monitored irrigation consumption on a portion of the xeric landscape. Submeters were typically read monthly, as with mainmeters and were used to study per-unit area application of water comparatively. The area monitored by the submeter was called the Xeric Study Area. Study areas were tied to irrigation zones and stations. Virtually all study properties have in-ground irrigation systems and controllers to avoid the presence or absence of these as a major confounding factor. This experimental control is important because it has been noted that the presence of automated irrigation is highly associated with increased water usage for residential properties (Mayer and DeOreo<sup>8</sup> et al. 1999) apparently because such systems make irrigation more likely to occur regularly versus hand-watering. Having participants in both groups possess automated systems also avoids the potential bias of more heavily turf-covered properties being more likely to be fully automated, thus having higher consumption as was the case for Bent<sup>1</sup> 1992 (as identified in Gregg<sup>4</sup> et al. 1994). All areas of each property were broken down into landscape categories. For example, a XS Group property might have monitored (via the submeter) xeric landscape and unmonitored xeric, turf, garden, and

other (non-landscaped) areas. Square footages were recorded for each of these respective area types.

In addition to water-consumption monitoring, residents agreed to a yearly site visit for data-collection purposes. During site visits, information was collected on the xeric species present, plant canopy coverage at the site, components of the irrigation system, and per-station flow rates.

Staff trained in the identification of locally used landscape plants collected data on plant size and species present.

Plant canopy coverage was calculated by first taking the observed plant diameters, dividing this number by two to get radius, then applying the formula for getting the area of a circle ( $A=\pi r^2$ ). This area result was then multiplied by the quantity of those species of plants observed to be at that size. The summation of all areas of all plants of all size classes in the study area is the total canopy coverage for that area.

Data on the components of irrigation systems was collected by staff trained in the different types of irrigation emitters available (ex. drip, microsprays, bubblers, etc.). Staff then ran individual stations and watched meter movement to get the per-station flow rates.

The Turf Study (TS) Group is composed of properties of more traditional landscape design, where an average 2,462 sqft of the landscaped area was of traditional turfgrass (most commonly Fescue). Mainmeter data was collected in the same manner as for the XS Group. Due to design challenges, the submeter was more commonly hooked to monitor a mixed type of landscape rather than just turf, though many did exclusively monitor turf (only “exclusively turf” monitoring configurations were used in per-unit area landscape analyses). TS participants enrolled voluntarily, without an incentive and agreed to yearly site visits as above. Other data on irrigation systems was collected in a manner similar to that for the XS Group properties. A total of 253 residences were recruited into the TS Group.

The enrollment of participant residences into the XS and TS Groups was directly dependent on homeowners’ willingness to participate in this study. For this reason, sampling bias was of reasonable concern to SNWA. To address this, a third subset of non-contacted Comparison Groups was created to evaluate potential biases. Comparison properties were properties with similar landscape footprints and of similar composition to the TS group and pre-conversion XS Group and were in the same neighborhoods as these treatment properties. This control group was also subject to the same water rates, weather, and conservation messaging as the treatment groups. Having this group also permitted SNWA to evaluate the combined effects of submetering and site visits on the treatment groups.

## **GENERAL DATA METHODS, STRATEGIES, AND STATISTICS**

Several different data analysis methods were applied in the course of the study. Details of each can be found in the corresponding subsections below. Broadly, analysis methods fell into the categories of pre- vs. post-treatment evaluations, comparative analyses of different treatment groups, analyses to determine variables associated with consumption, and assorted cost-benefit analyses. Statistical methods employed include descriptive statistics (ex. means, medians, etc.), tests for differences in means assuming both normally distributed data (*t*-tests) and non-normally distributed (i.e., non-parametric) data (*Mann-Whitney U*-tests), as well as techniques employing established economic principles and multivariate regression (some details of regression models are included in Appendix 2). In means comparisons, statistical significance was determined to occur when the probability of a Type I error was less than 5% ( $\alpha=0.05$ ). Presentation of data involving calculations of differences in values (for example, means differences) may not appear to add up in all cases, due to rounding. Types of data analyzed include mainmeter consumption data, submeter consumption data combined with area data (i.e., application per unit area data), flow-rate data, cost data, survey responses, and assorted demographic and Clark County Assessor's Office data. Consumption data was gathered by the aforementioned purveyor entities and assembled by SNWA. Most other data was collected by SNWA (Aquacraft Inc. also performed some analyses on consumption and data logger collected data under contract to SNWA). In many analyses, data was scatterplotted and objective or subjective outlier removal done as deemed appropriate. Finally, in some cases, data analysis was expanded upon to include attempts at modeling. These endeavors are elaborated on in other parts of the manuscript.

### **PRE/POST ANALYSES**

For each property and year where complete monthly consumption records were available, these were summed to provide yearly consumption. Data for each XS Group property was assembled from the five years before conversion (or as many records as were available; only properties having converted from turf to xeriscape were in this analysis sample) and as many years post-conversion as records permitted up through 2001. These data sets permitted comparison of total yearly consumption before and after the landscape conversion. The impact of submetering and site visits could also be evaluated by comparing mainmeter records for the TS Group pre- and post-installation of landscape submeters. Differences could be further confirmed by comparing the change in total household consumption following the conversion or submetering event for the XS and TS groups respectively against the change in consumption for non-contacted, non-retrofitted properties of similar landscape composition. The general analysis strategy for Objective 2 of the approved Scope (Appendix 1) is summarized in the following tables (Tables 1 and 2):

**TABLE 1: Planned Pre-/Post-Retrofit Analyses for XS Group**

	<b>Pre-retrofit (kgal/yr)</b>	<b>Post-retrofit (kgal/yr)</b>	<b>Difference in Means (kgal/yr)</b>
<b>Xeriscape Treatment</b>			
<b>Comparison</b>			
<b>Difference in Means (kgal/yr)</b>			

**TABLE 2: Planned Pre-/Post-Retrofit Analyses for TS Group**

	<b>Pre-retrofit (kgal/yr)</b>	<b>Post-retrofit (kgal/yr)</b>	<b>Difference in Means (kgal/yr)</b>
<b>Submetered Conventionally Landscaped Treatment</b>			
<b>Comparison</b>			
<b>Difference in Means (kgal/yr)</b>			

**ANALYSES OF SAVINGS OVER TIME AND SEASONS**

Objective 3 directs SNWA to measure the variability of water savings over time and across seasons. In the approved Scope, this was anticipated to involve comparing the XS, TS, and Comparison Groups to derive savings estimates in the manner specified in the tables that follow (Tables 3 and 4):

**TABLE 3: Planned Post-Retrofit Analyses for XS Group Across Time**

	<b>First Year's Consumption (Y1)</b>	<b>Third Year's Consumption (Y3)</b>	<b>Difference in Means (kgal/yr)</b>
<b>Xeriscape Treatment</b>			
<b>Comparison</b>			
<b>Difference in Means (kgal/yr)</b>			

**TABLE 4: Planned Post-Retrofit Analyses for TS Group Across Time**

	<b>First Year's Consumption (Y1)</b>	<b>Third Year's Consumption (Y3)</b>	<b>Difference in Means (kgal/yr)</b>
<b>Submetered Conventionally Landscaped Treatment</b>			
<b>Comparison</b>			
<b>Difference in Means (kgal/yr)</b>			

Since in most cases, meters were read monthly or at least bimonthly, SNWA is able to provide an analysis exceeding the level of detail originally specified in the Scope. Specifically, the longevity of savings from conversions for *each* year following the conversion could be evaluated, thus the following new table specifies the more in-depth level for the “over time” analyses called for in Objective 3:



**TABLE 5: Enhanced Post-Retrofit Analyses for XS Group Across Time**

<b>Mean Post-retrofit Consumption</b>	<b>First Year Post-retrofit (Y1)</b>	<b>Second Year Post-retrofit (Y2)</b>	<b>Third Year Post-retrofit (Y3)</b>	<b>Fourth Year Post-retrofit (Y4)</b>	<b>Fifth Year Post-retrofit (Y5)</b>
<b>Xeriscape Treatment (kgal/year)</b>					
<b>Comparison Group (kgal/year)</b>					
<b>Difference in Means (kgal/year)</b>					

**TABLE 6: Enhanced Post-Retrofit Analyses for TS Group Across Time**

<b>Mean Post-retrofit Consumption</b>	<b>First Year Post-retrofit (Y1)</b>	<b>Second Year Post-retrofit (Y2)</b>	<b>Third Year Post-retrofit (Y3)</b>	<b>Fourth Year Post-retrofit (Y4)</b>	<b>Fifth Year Post-retrofit (Y5)</b>
<b>Submetered Conventionally Landscaped Treatment (kgal/year)</b>					
<b>Comparison Group (kgal/year)</b>					
<b>Difference in Means (kgal/year)</b>					

Recruitment of properties for the XCS spanned a couple of years. For this reason, in order to evaluate true changes over time, the first year after each conversion was designated as Y1, the second as Y2, and so forth. As such, consumption data for a property starting in, for example, 1995, was designated as belonging to Y1, but for a different property starting in 1996, 1996 was Y1. In this way, the impact of different start years was corrected for and multiyear analyses could be considered on a more common basis. This permits inferences to be made about how landscape water consumption and savings change over time as plants in the xeric areas mature. It is also the reason the sample size appears to diminish for the XS Groups from Y1 to Y5. It is not that there was heavy loss of sample sites, rather that fewer sites were in existence for a total of five years owing to early enrollment. A similar effect is seen in the TS Group. There is no data for Y5 for the TS Group because enrollment for that Group started later than for the XS Group.

Savings from xeriscape may be greatest in summer when evapotranspirational demand is greatest for all plants, but so to an extreme degree in southern Nevada for turfgrasses (Source: University of Nevada Cooperative Extension). In considering how savings may be different across seasons, the Scope (Appendix 1) directs the SNWA to certain prescribed analyses (Tables 7 and 8):

**TABLE 7: Planned Summer Post-Retrofit Analyses for XS Group**

	<b>Pre-Retrofit Summer Consumption (kgal/month)</b>	<b>Post-Retrofit Summer Consumption (kgal/month)</b>	<b>Difference in Means (kgal/month)</b>
<b>Xeriscape Treatment</b>			
<b>Comparison Group</b>			
<b>Difference in Means (kgal/month)</b>			

**TABLE 8: Planned Summer Post-Retrofit Analyses for TS Group**

	<b>Pre-Retrofit Summer Consumption (kgal/month)</b>	<b>Post-Retrofit Summer Consumption (kgal/month)</b>	<b>Difference in Means (kgal/month)</b>
<b>Submetered Conventionally Landscaped Treatment</b>			
<b>Comparison Group</b>			
<b>Difference in Means (kgal/month)</b>			

Because of the resolution available by submetering, even more detailed data pertaining to application of water to turf and xeriscape through seasons is available in the comparative per-unit area irrigation analyses (see following section and Comparison of Per-Unit Area Water Application between Turfgrass and Xeric Landscape in *Results and Discussion*).

**COMPARATIVE PER-UNIT AREA IRRIGATION ANALYSES**

Submeter consumption data combined with measurement of the irrigated area permitted calculation of irrigation application on a per-unit area basis (gallons per square foot, which can also be expressed as precipitation inches equivalents) for most study participants. In this way, exacting measures of consumption for irrigation of xeric and turf landscape types could be measured. The sample size ( $N_s$ ) is the product of the number of months or years of data and the number of valid submeter records analyzed. Sample sizes for specific analyses appear in *Results and Discussion*. Only records for submeters that monitored turf exclusively were included in per-unit area analyses involving the TS Group so that other landscape types would not confound calculation of results.

No prescribed analyses of submeter consumption data appear in the Scope. The two basic sets of analyses selected by SNWA were (i.) a comparative analysis of annual application to xeric and turf areas and (ii.) a comparative analysis of monthly application to xeric and turf areas. The analytical setup of these appears in Tables 9 and 10 respectively. Secondary analyses comparing usage to theoretical reference ET demand projections follow the basic comparisons and appear in *Results and Discussion*.

**TABLE 9: Planned Comparative Analysis of Turf and Xeric Per Unit Area Annual Application**

	<b>Per Unit Area Application (gallons/square foot/year)</b>
<b>Submetered Turf (TS Group)</b>	
<b>Submetered Xeriscape (XS Group)</b>	
<b>Difference (gallons/square foot/year)</b>	

**TABLE 10: Planned Comparative Analysis of Turf and Xeric Per Unit Area Application for Each Month**

	<b>Jan Gal/SqFt</b>	<b>Feb Gal/SqFt</b>	<b>Mar Gal/SqFt</b>	<b>Apr Gal/SqFt</b>	<b>May Gal/SqFt</b>	<b>Jun Gal/SqFt</b>	<b>Jul Gal/SqFt</b>	<b>Aug Gal/SqFt</b>	<b>Sep Gal/SqFt</b>	<b>Oct Gal/SqFt</b>	<b>Nov Gal/SqFt</b>	<b>Dec Gal/SqFt</b>
<b>Submetered Turf (TS Group)</b>												
<b>Submetered Xeriscape (XS Group)</b>												
<b>Difference (gallons/square foot/month)</b>												

## MULTIVARIATE ANALYSES TO IDENTIFY SIGNIFICANT SOURCES OF VARIABILITY

Objective 4 of the Scope (Appendix 1) directs SNWA to assess variability of water use amongst the study participants and identify what factors contribute to that variability. Potential sources of variability originally specified for investigation in the Scope included the following:

- Number of members in the household
- Age of occupants
- Number of bathrooms
- Income
- Home value
- Percentage of xeriscaping
- Xeriscape density
- Turf type
- Type of irrigation
- Lot size
- Landscapeable area
- Existence of a pool
- Flow rates
- Water use factors

As the XCS developed, additional potential factors were assessed. A complete listing of data recorded is included in Appendix 3 (not all data was collected for all properties in the study).

Preliminary investigations focused on some of the above variables from a principally univariate analysis perspective (DeOreo<sup>9</sup> et al. 2000, Sovocool<sup>10</sup> et al. 2000, Sovocool and Rosales<sup>11</sup> 2001, Sovocool<sup>12</sup> 2002). The advantage of this was that it permitted rapid quantification and association of target variables' influences on participant water use, especially at the per-unit area scale. However, the most sophisticated way to deal with a study of this type where there are a number of potential independent associations of several predictor variables to a dependent variable is by the application of multivariate regression analysis methods. This permits so-called "partial regression" of independent variables to the target dependent one, here water consumption. Multiple regression for estimation can be expressed in the general multiple regression equation as follows:

$$\hat{Y}_i = \hat{a} + \hat{b}_1 * X_{1i} + \hat{b}_2 * X_{2i} + \dots + \hat{b}_{ni} * X_{ni} + \epsilon$$

Where  $\hat{Y}$  is the estimated dependent variable,  $\hat{a}$  is the y-axis intercept,  $\hat{b}$  is each estimated beta partial regression coefficient representing the independent contribution of each independent variables' influence on  $\hat{Y}$ ,  $X$  is each independent variable up to the  $n$ th variable,  $i$  is the time period and  $\epsilon$  is the error term for the model.

Multicollinearity between  $X$  variables violates the underlying assumptions of regression models and can be dealt with by setting limiting tolerance thresholds of similarity in contribution of variability to a regression model. This, in turn, permits identification and possible exclusion of such highly collinear and possibly inappropriate independent variables. The most significant variables can then be quantified and their relative vector and magnitude of association on the

dependent variable can be deduced, ultimately yielding an explanatory multivariate model of how such variables may contribute to water consumption. Such variables are explored for association to total household consumption and xeric landscape submeter consumption in the results section in two distinct modeling exercises.

## **ECONOMIC ANALYSES**

Objective 5 of the Scope mandates quantification and measurement of capital costs and maintenance costs of the conversion. In the summer of 2000, data on landscape maintenance economics was obtained via surveys sent to study participants. The survey helped quantify both labor hours and direct costs associated with landscape choices. For details on the survey and methodology, consult Hessling<sup>13</sup> (2001). Three hundred surveys were returned for analysis. Results of these were tabulated and compiled, and analyses proceeded from there.

By the very nature of the study methodology, it was recognized at the outset that a simple comparison of the XS and TS groups would likely fail to demonstrate the economic considerations with respect to maintenance of the whole landscape level as most residents' landscapes were composed of multiple landscape types (at the least, both xeric and turfgrass areas). This led to an analytical method of comparing the costs of landscape maintenance based on the relative percentages residents had of turf and xeric areas respectively.

The water bill savings associated with conversion projects were calculated based on the Las Vegas Valley Water District's water rates as they currently stand (in early 2004). Savings were calculated by modeling bills for a typical fifth decile (midrange in consumption) home where the average yearly consumption is 208,057 gallons and for such a home doing an average (according to data collected for the Water Smart Programs single-family sector in early 2004) 1,615.8-sqft-conversion from turfgrass to xeric landscape (note the difference in this average size conversion relative to that of the XS Study Group; conversion sizes, along with lot sizes, have diminished over time in this area). Bills were modeled on a monthly basis and all charges were applied that actually appear for customers. An example output of this model appears in Appendix 4.

As directed in the Scope (Appendix 1), the financial viability of xeriscape conversions was explored. This necessitated looking at the economics of conversions from the homeowner and SNWA perspectives. Hessling<sup>13</sup> (2001) attempted some of these initially. A follow-up analysis from these same perspectives was performed in the writing of this report and is included in *Results and Discussion*. The homeowner perspective included an estimative Net-Present-Value (NPV)-based modeling approach to determine when return on investment (ROI) was achieved and details on this model appear in Appendix 5. This same model is used to determine the incentive level necessary to induce change (Objective 6) by making some assumptions about what timeframe is acceptable for owners to achieve ROI. The approach used for the SNWA perspective is to consider alternative sources of water and use the cost associated with these to determine the maximum amount SNWA should pay to help convert grass to xeric landscape.

## Results and Discussion


### REDUCTION IN TOTAL HOUSEHOLD WATER CONSUMPTION FOLLOWING CONVERSION TO XERISCAPE

Results for the XS Group pre/post-conversion comparisons are shown in Table 11 and Figure 1.

**TABLE 11: Pre-/Post-Retrofit Analyses for XS Group**

	Pre-retrofit (kgal/yr)	Post-retrofit (kgal/yr)	Difference in Means (kgal/yr)	t-tests (* denotes significance)
<b>Xeriscape Treatment n=321</b>	Mean=319 Median=271	Mean=223 Median=174	<b>96*</b> (30% reduction from pre-retrofit)	<b>t=16.8*</b> <b>p&lt;0.01</b>
<b>Comparison n=288</b>	Mean=395 Median=315	Mean=382 Median=301	<b>13</b> (3% reduction from pre-submetering)	<b>t=1.85</b> <b>p=0.07</b>
<b>Difference in Means (kgal/yr)</b>	<b>76*</b>	<b>159*</b>		
<b>t-tests (* denotes significance)</b>	<b>t=4.32*</b> <b>p&lt;0.01</b>	<b>t=9.69*</b> <b>p&lt;0.01</b>		

FIGURE 1: Pre-/Post-Retrofit Consumption for XS and Comparison Groups



Mean monthly consumption for the residences dropped an average of 30% following conversion. A dependent *t*-test demonstrates that the reduction in usage is highly significant ( $t=16.8$ ;  $p<0.01$ ).

Though individual performance may vary greatly, the overwhelming majority of homes in the study saved water following the conversion (285 out of 321 analyzed). This finding of about a third reduction in consumption is nearly identical to findings from a study of residences in Mesa, Arizona (Testa and Newton<sup>2</sup> 1993). It may be that a reduction of about this percentage may be anticipated to occur when the average single-family residence built in the late 20<sup>th</sup> century does an average-size conversion in the southwestern United States. The large savings are likely in part because the great majority of water consumption goes to outdoor irrigation in this region. In this study, the average savings realized was 96,000 gallons per year per residence.

The difference in consumption of the pre-retrofit homes to the non-contacted comparison homes is shown in Table 11 and Figure 1. As demonstrated, a *t*-test of consumption between these two groups shows there was significant difference in initial consumption between the two groups (*t*=4.32; *p*<0.01), suggesting self-selection bias. This is not surprising since recruitment of study participants was voluntary. People who were already conserving more were apparently more likely to enroll and agree to convert a portion of their respective properties. This does not however invalidate the results, as (i.) this incentive-based approach is essentially the same as the approach used for enrolling people in the actual program SNWA has (see Appendix 5) and, more importantly (ii.), there is no compelling evidence that the Comparison Group experienced significant reduction over the same time period so the savings are likely attributable exclusively to the landscape conversion.

The analysis procedures in the Scope (Appendix 1) suggest that the impact of submetering on outdoor irrigation may be revealed by comparing consumption at the conventionally landscaped properties with submeters (the TS Group) to that for the associated comparisons for that Group. The data appearing in Table 12 fulfill this prescribed Scope treatment.

**TABLE 12: Pre-/Post-Retrofit Analyses for TS Group**

	<b>Pre-submetering (kgal/year)</b>	<b>Post-submetering (kgal/year)</b>	<b>Difference in Means (kgal/yr)</b>	<b>t-tests (* denotes significance)</b>
<b>Submetered Conventionally Landscaped Treatment n=205</b>	<b>Mean=352</b> Median=303	<b>Mean=319</b> Median=268	<b>34*</b> (10% reduction from pre-retrofit)	<b>t=5.08*</b> <b>p&lt;0.01</b>
<b>Comparison n=179</b>	<b>Mean=364</b> Median=314	<b>Mean=347</b> Median=296	<b>17*</b> (5% reduction over timeframe)	<b>t=2.08*</b> <b>p&lt;0.05</b>
DIFFERENCE IN MEANS (KGAL/YR)	<b>12</b>	<b>28</b>		
T-TESTS (* DENOTES SIGNIFICANCE)	<b>t=0.52</b> <b>p=0.60</b>	<b>t=1.41</b> <b>p=0.16</b>		



There are two potential issues though with trying to consider this analysis an evaluation of the effectiveness of submetering. First, submetering is typically studied where the scenario is one where water consumption through the submeter is relayed to end-use customers and where the customers are billed for it. Without consumption data and billing, the residents in this study have received no price signal to encourage them to read the meter or reduce consumption. This theory corresponds with what staff members have observed in the field with respect to the behavior of customers. Most participants apparently did not even think about the meter until it was time for their yearly site review and often they stated they had forgotten it was even there. So here, the dynamic of submetering is rather unique and the impact most likely minimal.

The second consideration, at least as potentially significant, is the fact that participants had been exposed to annual site visits, which is likely a more important variable in terms of modifying behavior (no conservation training or formal education took place at site visits, though staff members did answer questions posed to them). Indeed, the Comparison Group provides for a good gauge of the impacts on treatment groups due to site visits. Initially, results seem to suggest a reduction of possibly up to 34,000 gallons annually associated with visits and submetering ( $t=5.08$ ;  $p<0.01$ ) though, as revealed in the next analyses, this impact appears to be only temporary (seen only in the first year, Table 15) and is probably in actuality much more negligible given half the “reduction” also appears to have taken place in the control group ( $t=2.08$ ,  $p<0.05$ ). The control group reduction may be due to background conservation at the community level.

With respect to understanding how submetering with consumption billing may be of conservation benefit, a national research effort (Mayer et al. 2004<sup>14</sup>), supported in part by SNWA, has just been completed which provides much more insight into the benefits of submeters for water conservation purposes (also see Rosales<sup>15</sup> et al. 2002).


## **ASSESSMENT OF SAVINGS POTENTIAL ACROSS TIME AND SEASONS**

For the XS Group, significant reduction in total yearly consumption took place immediately following conversion and remained relatively stable at that decreased level through subsequent years, showing no erosion with time (Table 13 and Figure 2). In every year, the XS Group consistently had lower consumption than the Comparison Groups, and this was statistically significant (Table 13). This suggests that conversions are a viable way to gain substantial water savings over at least a medium-term timeframe and quite possibly over a long one as well. It also resolves questions about whether or not xericape takes more water in the first year following conversion (apparently the answer is no) and it suggests that, at least over the medium-term, there is no erosion of savings obtained from conversions due to residents’ response to growth of plants in their xeric areas.

For the XS Group, the relative reduction in consumption became even more pronounced in the summer (Table 14) where, savings averaged 13,000 gallons per summer month (Table 14:  $t=18.5$ ;  $p<0.01$ ) versus an average of 8,000 per month over the entire year. It should be noted that a very small, but statistically significant reduction of 1,600 gallons per month appears to have also taken place in the Comparison Group during the summer (in a pre- vs. post-comparison of the study timeframe, Table 14:  $t=1.98$ ;  $p<0.05$ ). Overall, the results are consistent with the theory that xeric landscapes save the most during the summer. The comparative per-unit analyses that follow reveal why this is the case.

In considering savings stability over extended time, it was found that the submetered TS group only demonstrated significantly decreased consumption for the first year following retrofit, after which savings were not significant (Table 15; statistics in table). This initial reduction might be due to residents' interest in the research and in conservation when new to the study, this wearing off with time. Again, it is important to recall that in no single year was the consumption statistically different from the comparison group properties. The submetered TS Group did have significantly lower consumption in the summer, with a savings of 3,300 gallons per month (Table 16:  $t=3.78;p<0.01$ ) whereas the comparison group to the TS Group showed no such reduction (Table 16:  $t=1.03;p=0.31$ ). However, there was no difference in average monthly summer consumption between the submetered properties and the controls after the retrofit (Table 16:  $t=1.03;p=0.31$ ). Overall the results in Table 15 seem to reflect the finding that little enduring change in consumption was achieved by the TS Group over time despite submeter installation.

**FIGURE 2: Pre-/Post-Retrofit Consumption for XS Group Across Time**



**TABLE 13: Enhanced Post-Retrofit Analyses for XS Group Across Time**

<b>Post-retrofit Consumption</b>	<b>First Year Post-retrofit (Y1)</b>	<b>Second Year Post-retrofit (Y2)</b>	<b>Third Year Post-retrofit (Y3)</b>	<b>Fourth Year Post-retrofit (Y4)</b>	<b>Fifth Year Post-retrofit (Y5)</b>
<b>Xeriscape Treatment (kgal/year)</b>	<b>214<sup>Δ</sup></b> (32% reduction from pre-retrofit) n=320	<b>220<sup>Δ</sup></b> (30% reduction from pre-retrofit) n=318	<b>227<sup>Δ</sup></b> (28% reduction from pre-retrofit) n=306	<b>211<sup>Δ</sup></b> (33% reduction from pre-retrofit) n=211	<b>202<sup>Δ</sup></b> (36% reduction from pre-retrofit) n=61
<b>Comparison Group (kgal/year)</b>	<b>372</b> n=280	<b>387</b> n=275	<b>383</b> n=260	<b>362</b> n=183	<b>345</b> n=54
<b>Difference in Means (kgal/year)</b>	<b>158</b>	<b>167</b>	<b>156</b>	<b>151</b>	<b>143</b>
<b>t-tests (* denotes significance)</b>	<b>t=9.98*</b> <b>p&lt;0.01</b>	<b>t=9.29*</b> <b>p&lt;0.01</b>	<b>t=9.08*</b> <b>p&lt;0.01</b>	<b>t=8.02*</b> <b>p&lt;0.01</b>	<b>t=4.85*</b> <b>p&lt;0.01</b>

Treatment group values with a <sup>Δ</sup> are significantly lower than pre-retrofit value.

**TABLE 14: Summer Post-Retrofit Analyses for XS Group**

	<b>Pre-Retrofit Summer Consumption (kgal/month)</b>	<b>Post-Retrofit Summer Consumption (kgal/month)</b>	<b>Difference in Means (kgal/month)</b>	<b>t-tests (* denotes significance)</b>
<b>Xeriscape Treatment n=321</b>	Mean=38 Median=31	Mean=25 Median=19	<b>13*</b>	<b>t=18.5* p&lt;0.01</b>
<b>Comparison Group n=288</b>	Mean=47 Median=38	Mean=46 Median=35	<b>1.6*</b>	<b>t=1.98* p&lt;0.05</b>
<b>Difference in Means (kgal/month)</b>	<b>9*</b>	<b>21*</b>		
<b>t-tests (* denotes significance)</b>	<b>t=4.23* p&lt;0.01</b>	<b>t=10.1* p&lt;0.01</b>		

**TABLE 15: Enhanced Post-Retrofit Analyses for TS Group Across Time**

<b>Post-submetering Consumption</b>	<b>First Year Post-submetering (Y1)</b>	<b>Second Year Post-submetering (Y2)</b>	<b>Third Year Post-submetering (Y3)</b>	<b>Fourth Year Post-submetering (Y4)</b>	<b>Fifth Year Post-submetering (Y5)</b>
<b>Submetered Conventionally Landscaped Treatment (kgal/year)</b>	<b>291<sup>Δ</sup></b> (6% decrease from pre-submetering) n=228	<b>312</b> (1% increase from pre-submetering) n=229	<b>317</b> (2% increase from pre-submetering) n=228	<b>315</b> (2% increase from pre-submetering) n=146	<b>No Data Available</b>
<b>Comparison Group (kgal/year)</b>	<b>332</b> n=170	<b>357</b> n=173	<b>351</b> n=167	<b>351</b> n=108	<b>No Data Available</b>
<b>Difference in Means</b>	<b>41</b>	<b>45</b>	<b>34</b>	<b>36</b>	
<b>t-tests (* denotes significance)</b>	<b>t=2.28</b> <b>p=0.02</b>	<b>t=2.39</b> <b>p=0.02</b>	<b>t=1.65</b> <b>p=0.10</b>	<b>t=1.40</b> <b>p=0.16</b>	

Treatment group values with a <sup>Δ</sup> are significantly lower than pre-submetering value.

**TABLE 16: Summer Post-Retrofit Analyses for TS Group**

	<b>Pre-Submetering Summer Consumption (kgal/month)</b>	<b>Post-Submetering Summer Consumption (kgal/month)</b>	<b>Difference in Means (kgal/month)</b>	<b>t-tests (* denotes significance)</b>
<b>Submetered Conventionally Landscaped Treatment n= 205</b>	Mean=41.7 Median=34.0	Mean=38.5 Median=31.0	3.3*	t=3.78* p<0.01
<b>Comparison Group n=179</b>	Mean=42.0 Median=36.0	Mean=41.0 Median=34.7	1.0	t=1.02 p=0.31
<b>Difference in Means (kgal/month)</b>	0.3	2.5		
<b>t-tests (* denotes significance)</b>	t=0.97 p=0.92	t=1.03 p=0.31		

**COMPARISON OF PER-UNIT AREA WATER APPLICATION BETWEEN TURFGRASS AND XERIC LANDSCAPE**

**Annual application**

Annual per unit area irrigation application data summaries are found in Table 17 and Figures 3 and 4. There was a great difference in the annual water application to turf and xeric landscape areas (Table 17 and Figure 3). Turf received an average of 73.0 gallons per square foot annually (117.2 inches), while xeriscape received on average, just 17.2 gallons (27.6 inches) each year (only 23.6% of the amount of water applied for turfgrass maintenance). The difference was thus 55.8 gallons per square foot per year (89.6 inches), and this was found to be highly significant assuming a normal distribution of data ( $t=27.0$ ;  $p<0.01$ ).


**TABLE 17: Annual Per-Unit Area Application to Turf and Xeriscape**

	<b>Per Unit Area Application (gallons/square foot/year)</b>	<b>Per Unit Area Application (inches/year)</b>	<b>Sample Distribution Statistics</b>
<b>Submetered Turf (TS Group) <math>n_s=107</math></b>	<b>Mean=73.0</b> Median=64.3	<b>Mean=117.2</b> Median=103.2	<b>Standard Deviation=40.0</b> <b>Skewness=1.17</b> <b>Kurtosis=1.36</b>
<b>Submetered Xeriscape (XS Group) <math>n_s=1550</math></b>	<b>Mean=17.2</b> Median=11.5	<b>Mean=27.6</b> Median=18.5	<b>Standard Deviation=18.6</b> <b>Skewness=3.14</b> <b>Kurtosis=14.9</b>
<b>Difference (gallons/square foot/year)</b>	<b>Mean=55.8</b>	<b>Mean=89.6</b>	
<b>t-tests (* denotes significance)</b>	<b><math>t=27.0^*</math></b> <b><math>p&lt;0.01</math></b>		
<b>Levene's Test (* denotes significance)</b>	<b><math>F(1, 1655)=130.3^*</math></b> <b><math>p&lt;0.01</math></b>		
<b>Mann-Whitney U Test (* denotes significance)</b>	<b><math>U=10177</math></b> <b><math>z=15.2^*</math></b> <b><math>p&lt;0.01</math></b>		

Detailed statistics were not generated for the small set of multifamily and commercial sites; however, the average consumption on those xeric areas where viable data could be collected was 16.7 gallons per square foot per year ( $n_s=22$ ). This suggests the use of xeric landscape in these sectors may result in similar savings as that observed above on a comparative landscape basis

(i.e., savings of ca. 55.8 gallons per square foot annually versus what application would have been for turf).


**FIGURE 3: Annual Per Unit Area Application to Turf and Xeriscape**



Distinct differences in the sample distributions for the XS and TS irrigation data were of concern from a statistical analysis perspective. Both distributions had features strongly suggesting data was not distributed homogeneously across the two groups (Table 17 and Figure 4). In particular, the XS Group data was heavily skewed with the vast majority of participants using very little water. Turf application, while indeed skewed, appears almost normal compared to xeric application, which is very heavily skewed (skewness = 3.14) and peaks sharply (kurtosis=14.9) at the lower end of the distribution. This is because the vast majority of XS participants used a *very* small amount of water to irrigate their xeric areas, while a handful used greatly more volume on theirs. Because *t*-tests assume normality, the atypical and non-congruent distributions were of sufficient concern to justify running a Levene’s Test simultaneous with the *t*-tests to assess the potential need to apply non-parametric analytical techniques (though in practice the need for normality is lessened with large sample sizes due to the tendency of such a collection of data to mimic a normal distribution; aka. the central limit theorem). Indeed, the Levene’s Tests demonstrated significant differences in the distributions [Levene  $F(1,1655) = 130.3; p < 0.01$ ]. This suggested the need to backup the findings with non-parametric approaches. *Mann-Whitney U* (a summation and ranking based approach to the problem) was chosen as a good backup test. Associated *z* statistics for this test with corresponding probabilities are thus reported with the results in Table 17 as supporting evidence for statistical difference in irrigation application between the groups.



**FIGURE 4: Distribution of Annual Per Unit Area Application Data for Turf and Xeriscape**



**Monthly Application**

Monthly submeter data summaries for the XS Group and exclusively monitored turf TS Group participants appear in Table 18. It should be noted that at times the interval between reads stretched over more than one month and thus the dataset for the monthly data is slightly different than that for the above annual comparison as only consumption data deemed complete and assignable to a given month could be included (sometimes consumption across a two-month gap was averaged to fill the gap). There were issues with resolution in monitoring because typically at least a thousand gallons had to pass through the meter between reads in order for the consumption figure to be advanced and registered by the reader, and sometimes this did not happen for XS Group submeters monitoring relatively small areas due to low consumption. Both these factors likely result in slight inflation of monthly consumption values for both groups and this indeed appears to be manifest if monthly averages are summed across the year (i.e., this per unit area consumption figure is slightly higher than the annual one calculated in the previous section). Still, on a monthly basis the data is generally valid and valuable in comparative analyses and in comparing water application to irrigation requirements. Per-unit area application data is displayed graphically in Figure 5.


**TABLE 18: Monthly Per-Unit Area Application to Turf and Xeriscape**

	<b>Jan</b> Gal/SqFt	<b>Feb</b> Gal/SqFt	<b>Mar</b> Gal/SqFt	<b>Apr</b> Gal/SqFt	<b>May</b> Gal/SqFt	<b>Jun</b> Gal/SqFt	<b>Jul</b> Gal/SqFt	<b>Aug</b> Gal/SqFt	<b>Sep</b> Gal/SqFt	<b>Oct</b> Gal/SqFt	<b>Nov</b> Gal/SqFt	<b>Dec</b> Gal/SqFt
<b>Submetered Turf (TS Group)</b>	<b>2.97</b>	<b>2.96</b>	<b>3.44</b>	<b>6.07</b>	<b>9.37</b>	<b>10.79</b>	<b>11.86</b>	<b>10.23</b>	<b>8.47</b>	<b>6.20</b>	<b>4.37</b>	<b>2.47</b>
	2.11	2.06	3.29	4.85	7.86	9.38	10.50	8.71	7.15	5.29	3.50	1.96
	n <sub>s</sub> =85	n <sub>s</sub> =85	n <sub>s</sub> =85	n <sub>s</sub> =88	n <sub>s</sub> =93	n <sub>s</sub> =93	n <sub>s</sub> =95	n <sub>s</sub> =96	n <sub>s</sub> =99	n <sub>s</sub> =105	n <sub>s</sub> =107	n <sub>s</sub> =106
<b>Submetered Xeriscape (XS Group)</b>	<b>1.16</b>	<b>0.87</b>	<b>0.99</b>	<b>1.43</b>	<b>1.64</b>	<b>2.01</b>	<b>2.24</b>	<b>2.27</b>	<b>2.22</b>	<b>1.66</b>	<b>1.35</b>	<b>0.91</b>
	0.46	0.43	0.57	0.83	1.08	1.30	1.40	1.39	1.27	1.02	0.77	0.48
	n <sub>s</sub> =1291	n <sub>s</sub> =1337	n <sub>s</sub> =1377	n <sub>s</sub> =1409	n <sub>s</sub> =1412	n <sub>s</sub> =1421	n <sub>s</sub> =1431	n <sub>s</sub> =1456	n <sub>s</sub> =1496	n <sub>s</sub> =1519	n <sub>s</sub> =1534	n <sub>s</sub> =1534
<b>Difference (Gallons/Sqft)</b>	<b>1.81</b>	<b>2.09</b>	<b>2.45</b>	<b>4.64</b>	<b>7.74</b>	<b>8.78</b>	<b>9.62</b>	<b>7.96</b>	<b>6.25</b>	<b>4.54</b>	<b>3.02</b>	<b>1.56</b>
<b>t-tests (* denotes significance)</b>	<b>t=73.36*</b> <b>p&lt;0.01</b>	<b>t=7.52*</b> <b>p&lt;0.01</b>	<b>t=13.33*</b> <b>p&lt;0.01</b>	<b>t=9.92*</b> <b>p&lt;0.01</b>	<b>t=29.87*</b> <b>p&lt;0.01</b>	<b>t=27.7*</b> <b>p&lt;0.01</b>	<b>t=26.22*</b> <b>p&lt;0.01</b>	<b>t=21.96*</b> <b>p&lt;0.01</b>	<b>t=13.15*</b> <b>p&lt;0.01</b>	<b>t=17.59*</b> <b>p&lt;0.01</b>	<b>t=13.45*</b> <b>p&lt;0.01</b>	<b>t=9.39*</b> <b>p&lt;0.01</b>
<b>Mann-Whitney U Tests (* denotes significance)</b>	<b>U=23499</b> <b>z=8.84*</b> <b>p&lt;0.01</b>	<b>U=18127</b> <b>z=10.54*</b> <b>p&lt;0.01</b>	<b>U=15959</b> <b>z=11.27*</b> <b>p&lt;0.01</b>	<b>U=14225</b> <b>z=12.14*</b> <b>p&lt;0.01</b>	<b>U=6824</b> <b>z=14.49*</b> <b>p&lt;0.01</b>	<b>U=4415</b> <b>z=15.10*</b> <b>p&lt;0.01</b>	<b>U=6062</b> <b>z=14.89*</b> <b>p&lt;0.01</b>	<b>U=9776</b> <b>z=14.13*</b> <b>p&lt;0.01</b>	<b>U=12307</b> <b>z=13.91*</b> <b>p&lt;0.01</b>	<b>U=14501</b> <b>z=14.04*</b> <b>p&lt;0.01</b>	<b>U=25290</b> <b>z=11.98*</b> <b>p&lt;0.01</b>	<b>U=31202</b> <b>z=10.62*</b> <b>p&lt;0.01</b>

Note: bold gal/sqft values are means; regular font gal/sqft values are medians

The first, most obvious finding from the graph is that, turf application exceeds xeric application by a large statistically significant margin in every month. Ultimately, this is what constitutes the large annual savings seen at the annual landscape application and total home consumption levels.

**FIGURE 5: Monthly Per-Unit Area Application for Turf and Xeric Areas**




The data also suggests, among other things, that the reason for the aforementioned enhancement of savings during the summer is because turf application peaks drastically in the summer whereas application to xeriscape does not. A graph of the difference between the groups (Figure 6) demonstrates this is the case, and the observed pattern in savings obtained each month parallels the pattern observed for turfgrass application (Figure 5). It appears that the reason xeriscape saves so much water in this climate is related as much to the high demand of turfgrasses vs. plantings of most other taxa as it is to any inherent aspect of xeric landscape *per se*. Furthermore, inefficiencies in spray irrigation system design, installation, and operation further contribute to the savings of having xeric landscape in place of turf because these inefficiencies even further drive up application to the turfgrass to the point that it is much higher than the rate of evapotranspiration over the same timeframe (Figure 7).

Additional inferences can be made about the application of water to turfgrass areas by the participants. Specifically, on average, whereas they irrigated relatively efficiently in the spring, with the onset of summer temperatures in May, residents quickly increased their application, ultimately going way above  $ET_0$ . Moreover, they tended to stay well above  $ET_0$  through November. While it is expected that due to system inefficiencies, a high  $K_c$  for Fescue (Source: Cooperative Extension Office), leaching fraction considerations, and other factors, application usually would tend to exceed  $ET_0$  for turfgrass locally, the pattern suggests that


overall people irrigate relatively efficiently in spring as the weather warms and  $ET_o$  rises, probably due to the immediate feedback they receive as the grass yellows in response to moisture deficits. As they observe their landscape beginning to show visible signs of stress due to deficit irrigation, they increase their application accordingly. However, in May, they appear to start overreacting to the increasing stress and increase irrigation to well over the requirement. In fall, they do not however appear to respond in a correspondent way “coming down the curve,” probably because they do not have the same sort of visual feedback mechanism as they do in spring (i.e., they do not view the grass being “too green,” wet, nor the occurrence of runoff as something amiss). The result is a long lag in returning to application rates more closely approximating  $ET_o$  in the fall and early winter (Figure 7).

**FIGURE 6: Monthly Per-Unit Area Savings (Turf Area Application– Xeric Area Application)**




It is more difficult to make similar types of inferences with respect to xeric area application. While there is research under way on a variety of desert taxa to attempt to quantify irrigation demand and there have been generalized attempts to model or approximate xeriscape need based on observations and fractions of reference  $ET_o$ , at this time it would be risky to make highly specific inferences. The relative flatness of the xeric curve in Figure 5 does though seem to suggest that residents may irrigate xeric areas inefficiently as they seem to show little response to demands of different seasons.

**FIGURE 7: Monthly Per-Unit Area Application to Turf and Reference Evapotranspirational Demand**




**FIGURE 8: Monthly Per-Unit Area Application to Xeric Areas and 1/3 of Reference Evapotranspirational Demand**



If one does assume a sometimes-used local “rule-of-thumb” which states that xeriscape requires about a third of what turf needs, one can compare per-unit area application for xeriscape and this modified reference value (Figure 8). Using a one-third  $ET_0$  value is not out-of-line with modification approaches employed by the Irrigation Association<sup>16</sup> (2001) or WUCOLS<sup>17</sup> (2000) for estimating needs of low-water-use woody taxa in high-temperature southwestern regions. It is quite noteworthy that the summation of monthly xeric-area application values yields a yearly xeric-area application usage of 30.1 inches per year - nearly identical to the summation of monthly  $.33(ET_0)$  values, which is 30.5 inches. This would appear, initially at least, to suggest that this rule of thumb may work quite well on average for approximating xeric landscape usage over broad spatial and long temporal scales, even if it may not precisely work in a given month.

Normalizing these aforementioned potential reference values and the absolute departure from these in observed water application may reveal insights about when during the year the greatest absolute potential savings can be obtained. In Figure 9, this is done such that the absolute difference between mean application and respective references is quantified and displayed. Here, “0” (reference) is  $ET_0$  for turf and  $.33(ET_0)$  for xeric landscape respectively.

**FIGURE 9: Absolute Departure in Irrigation Application from Derived Respective Reference  $ET_0$  Values (Turf and Xeric Areas)**



Even with the xeric reference but a third of  $ET_0$ , it appears that, in addition to the differences due to plant usage, much more water is wasted in application to turfgrass than to xeric landscape. The

greatest waste for turfgrass occurs in the period of May through November. Thus, any improvements in turfgrass irrigation efficiency during this timeframe will have the greatest absolute impact in terms of water conservation. Interestingly, the greatest absolute potential for savings for xeric areas is not during this period, but rather from September thru January. Indeed to look upon the graph, one might initially conclude that residents under-irrigate xeric areas in spring and summer. Caution should be observed though in this type of reasoning as the  $.33(ET_0)$  reference is only theoretical and developed here as a guideline. That stated, the findings may suggest that, on average, little potential exists during the spring and summer for significant water savings by irrigation improvements to xeriscape. Finally, on an absolute basis, little total potential appears to exist for squeezing additional conservation out of xeric landscapes as, considered over the span of an entire year, xeric area irrigation appears to be efficient.

In contrast, opportunities to save great volumes of water appear to exist for turf areas throughout most of the year. Significant overwatering appears to occur May through November; efficiency improvements will yield the most absolute benefit during this period of the year. But how does the issue appear when one considers the problem through the perspective of *when can the most readily obtainable savings be achieved?*

Considering absolute irrigation departure from reference as above gives insights into the total potential to save water through a variety of irrigation improvements. However, there is also the question of how much water could be saved principally by relatively simple improvements in *controller management*. Figure 10 is such an attempt to view the problem through this framework, where the blue line is  $ET_0$  for turf and  $.33(ET_0)$  for xeric areas respectively, and is equivalent to 100% of each respective types reference value or “perfect efficiency.” Absolute values for inches application were normalized by converting them to percent departure from normalized respective reference values. In this way the relative departure from these aforementioned references is displayed as a percent value.

**FIGURE 10: Relative Departure in Irrigation Application from Derived Respective Reference  $ET_0$  Values (Turf and Xeric Areas)**





Figure 10 may suggest that there are specific times of the year when people irrigate both turf and xeric landscapes more or less efficiency than the ideal. As interpreted from Figure 10, the most inefficient irrigation, in a relative sense, may actually occur during non-peak months if efficiency is defined to be the difference between theoretical requirement and application. Expanding on this type of analysis and breaking the above relative departure values into efficiency classes yielded a summary of when people appear to irrigate most and least efficiently (Figure 11).

**FIGURE 11: Relative Departure in Irrigation Application from Derived Respective Reference ET<sub>0</sub> Values (Turf and Xeric Areas)**



It is well understood that, in practice, there is no such thing as a perfectly efficient irrigation system and, for this reason, the green designation in Figure 11 includes relative applications ranging from subreference values to those up to 20% above reference (this allows that there is typically a need in practice to compensate for lacking distribution uniformity in irrigation systems).

Interpretation of Figure 11 suggests that both xeric and turf areas are irrigated relatively efficiently in the spring. Irrigation efficiency for turfgrass areas starts to decline in May to the point where significant waste starts to occur and this continues until about September. In contrast xeric irrigation continues to be quite efficient during this time. Around September, turf is starting to be very inefficiently watered, in a relative sense, owing to residents’ failure to respond to the lower rate of evapotranspiration and decrease irrigation accordingly. A similar, if less severe, pattern is observed for xeric area irrigation, where at this time, these areas are also beginning to be irrigated inefficiently, probably for the same reason. By November, both xeric and turfgrass areas are, on average, being severely over-irrigated and this pattern continues through the cool season until February. Finally, efficiency starts to recover and both areas are actually being irrigated under suggested reference values by the end of March.

It needs to be acknowledged that some of this conclusion includes theoretical and speculative reasoning, especially considering the lack of data on xeric landscape water requirements and the fact that in actuality stress impacts, including those from water stress, lag in woody vegetation (Kozlowski et al. 1990<sup>18</sup>) so efficiency as considered here is much harder to gauge. Nevertheless, again, failure of residents to more effectively tie controller management (irrigation frequency and duration) to the changing environmental conditions appears to be one of the most pressing reasons for efficiency losses in both study groups, it is just to a lesser extent (and much lesser absolute impact in gallons) for those with more xeriscape.



This set of analyses provides SNWA with quantitative data on what parts of the year it should focus its strongest controller-management-oriented conservation messaging. This could be considered the “low hanging fruit” in terms of water conservation; it is where messaging to effect changes that may not require significant work and monetary investments on the part of residents may produce significant water conservation results. To recap, the findings in this section suggest the most value can be obtained by targeting controller-management messaging to the late summer and early fall as people begin to depart from “reasonable” efficiency values owing to their collective failure to adjust irrigation down for the cooler, low ET season. Reemphasis of this messaging should continue all winter long.

The exploration of application per-unit area vs. reference values is important for making inferences about management efficiency of water application. This; however, should not obscure the result that on average, per-unit area, xeric landscapes in this study received much less water in totality (Figures 3 and 4) and the pattern of received irrigation showed much less tendency towards “peaking” (Figure 5) than those areas planted with turf.

#### **SOURCES OF SIGNIFICANT VARIABILITY IN SINGLE-FAMILY RESIDENTIAL CONSUMPTION**

As explained in Methodology, multivariate regression analyses were employed to identify and quantify sources of variability of mainmeter and xeric submeter data. Specifically, variables in the combined study groups are explored for association to total household consumption and, for the XS Group, to xeric landscape submeter consumption. Regression modeling proceeded with the goal being to yield an optimum combination of the highest reasonable R-squared value with due consideration given to maximizing the degree to which the model was “complete” (to the extent possible given the available collected data). Details of the final selected multivariate regression models appear in Appendix 2. Explanation and discussion of each variable included follow for each of the respective models.

Presented models are only designed to broadly assess variables’ impacts. The models presented here are “estimation” models as defined (see *Methodology*). These models are not intended for use as “engineering” or “computational” type model applications whereby collecting certain data one could be reasonably certain that the answer yielded would closely approximate the real consumption at a given property.

## Variability in Annual Residential Consumption

Discussions of the selected independent variables included in the annual consumption model for the dependent variable *annual residential consumption* (labeled MAINMETE) follow. Overall, the annual consumption model appears to be a very good “fit” (adjusted  $R^2=0.80$ ) for this type of work (Nelson<sup>3</sup> 1994, Gregg<sup>4</sup> et al. 1994, Gregg<sup>19</sup> et al. 1999). This is likely due as much to the strong tie between outdoor usage (and the ability of independent variables associated with outdoor use to be practically measured) as to any design elements or analytical methods associated with the study. While relatively strong for the sample size, it must be stressed that this model’s utility is mostly in terms of helping to uncover and, to some extent, explain variables discreet associations with consumption at single-family residences. Quantifications of these associations in the multivariate context are limited to only those variables deemed significant.

### TOTALTUR

#### Definition of Variable:

*The total amount of turf at a residence in square feet as determined by research personnel.* This includes all turf regardless of whether it is part of a submetered area and regardless of what type of grass it is.

#### Results and Discussion:

This was the most significant variable by far ( $t=14.86$ ), and was found to be strongly positively associated with single-family residential consumption. It is a principal component of the model, contributing the bulk of its strength ( $\beta=0.622$ ). The results suggest that consumption increases roughly 59.1 gallons annually for each square foot of turf at the average home. It then increases *further* if the grass is Fescue (the impact of Fescue vs. other grasses is further explored below). Since the alternative grass is almost always Bermuda, the result suggests the average application rate for this warm-season grass by the study participants is about 59 gallons per square foot (see variable FESCUE for more discussion on this).

It should be noted that earlier multivariate work attempted to deduce the influence of landscape type by scrutinizing how much xeric landscape was found at a residence (DeOreo<sup>8</sup> et al 2000). While this is an acceptable approach, the amount of turfgrass present appears to be much more closely correlated with total annual consumption and, when included, typically displaces xeric area as a significant variable in the final models developed. Furthermore, since the amount of xeriscape was not significant in multivariate context (nor were other individual landscape types) it should be understood that the savings developed by SNWA’s Water Smart Landscapes program are mostly due to it, in essence, being a turf-removal program more than an alternative-landscape-promotion program. The results also suggest further significant lowering of household consumption probably would not be yielded by permitting the owner to get a rebate for turf removal at the expense of a quality landscape (for example, incentivizing the aforementioned “zeroscapes” at a higher SNWA incentive rate since they have no vegetation and theoretically require no water – this has been suggested by some). Since the xeric area contribution to annual consumption is so small, the substantial loss in quality of life yielded for the small gains in

conservation realized by effectively hardscaping landscape areas makes the argument for choosing hardscape in place of xeriscape for water conservation a position difficult to defend.

### TOTVAL

Definition of Variable:

*The dollar value of the single-family residential study property as specified in the Clark County Assessor's Office database.* This should not be considered to equate to a home's market value.

Results and Significance:

The assessed monetary value of the property, like the amount of turf at a residence, was a very highly significant variable in the model ( $t=5.45$ ). It is reasonable to assume that higher value properties are associated with higher consumption because (i.) they are likely to contain larger homes with typically larger, possibly more extravagant water-intensive landscapes and (ii.) they are, by nature, likely to be inhabited by people of greater wealth who are less sensitive to the price of water and thus more likely to use a greater volume of it. In a multivariate context, annual water consumption on average increases ca. 2.1 gallons alongside each dollar increase in Assessor's Office property value.

That increased wealth is associated with greater individual consumption is a well-understood tenant of economics and is a well-established concept in understanding persons' household utility consumption patterns. The impact of wealth in a similar context was explored by Gregg<sup>19</sup> et al. (1999) where the impact of neighborhood wealth was a significant factor in determining water usage.

### NLTHOMEA

Definition of Variable:

*The age of the residence is calculated as the difference between the analysis year (2004) and the year of construction as recorded in the Clark County Assessor's Office database.* This should not automatically be taken to be the age of the landscape or even, necessarily, the exact age of the specific study residence due to the way many developments are built as components of phases in this community.

Results and Significance:

This was a quite significant variable ( $t=2.67$ ) and one easily worthy of inclusion in the model. On average, consumption increased ca. 1600 gallons for each additional year older the property was.

There are several potential reasons for this. First, older properties in the Las Vegas area tend, on average, to be larger and the ratio of hardscape footprint to landscapeable area is lower. Next, older properties are more likely to incorporate landscape elements heavy on traditional themes (i.e., large areas of turfgrasses) in contrast to newer residences with landscapes built in a time where water conservation began to be a significant consideration (in the 1990s restrictions on the amount of turfgrass that could be installed at single-family residences were passed). Older properties are more likely to have irrigation systems that incorporate lower-efficiency devices and

fixtures (ex. brass spray heads). Finally, as irrigation systems age they inevitably become less efficient and more likely to leak.

Aspects of indoor use also likely contribute to the pattern. The installation of high-efficiency, low-flow fixtures and appliances after being legally mandated is anticipated to have contributed to newer properties having, on average, lower consumption. Also, as fixtures wear they may leak for some time without notice (toilet flappers for example) so, without timely maintenance, older properties are more likely to have continuous indoor leaks further contributing to higher consumption. The increased efficiency gains in homes with newer fixtures have been well documented (see Mayer and DeOreo<sup>8</sup> et al. 1999) and the overall finding that older homes tend to have higher water consumption is not surprising.

### APROXINC

#### Definition of Variable:

*Approximate total household income as revealed by 2001 survey data.* To make the income survey question less intimidating, and more likely to generate valid, significant numbers of responses, the potential answers were categorical with ranges and it was explicitly stated that this question was optional. Analysis proceeded based on the mean values of response ranges. While a great number of participants did respond, many of course did not and income is, unsurprisingly, the most limiting of independent variables in the multiple regression.

#### Results and Significance:

It is to be expected that, everything else being equal, increasing household income would on average be associable with higher per-household consumption of all commodities. This is the case for water as well in this multivariate model, which suggests that, on average, annual consumption may increase on average ca. 3000 gallons for every \$10,000 rise in income level ( $t=2.16$ ). Some may be surprised this should be given the fact that indoor water use is relatively constant per capita across a range of conditions and thus the sensitivity of the relationship between water consumption and price is usually considered to be rather muted. But, while water is indeed inelastic by common economic standards, in the Southwest, where a high proportion is used outdoors, it may be considered to be more discretionary in nature, especially when that outdoor use is for irrigation of landscapes (instead of crops), which are after all just ornamental. Certainly this study suggests that income is an important consideration in water consumption, as have others. Furthermore, higher incomes could be considered to be well correlated with large houses, large landscapeable areas, and more lush landscapes, all of which further drive up consumption in their own right.

There was considerable discussion between the principal author and some reviewers as to whether or not the income data should be included in the model. The arguments for inclusion were that it was found to be a significant variable in most comparisons, it is a different indicator than home value in that the former is more indicative of wealth and the latter is more indicative of actual disposable income (which could be spent on water use beyond necessity), and that removing it significantly weakens the model. The arguments for removing it include the supposition that often people give erroneous or fictional answers to questions about income, that income is potentially highly covariate with home value, that home value is really a better proxy variable for

income (and indeed in many studies using multiple regression it has been used for this purpose), and that its deletion does not weaken models such as this. Finally significant improvement in model sample size would be obtained by removing income as many people opted not to report it and thus it is very limiting to the model's available degrees of freedom.

The author considered the arguments for and against inclusion of income data carefully and proceeded to investigate the relationship between income and home value. The results of a correlation analysis between these two variables showed relatively little correlation ( $R^2 = 0.288$ ) as did a scatterplot of the data. Nonetheless, the concern was valid enough (and the possibility of significantly more degrees of freedom of sufficient interest) to justify creation of an incarnation of the model without income as an independent model variable. This exercise however resulted in an increase in the standard error of the estimate (i.e., an increased error of over 7,000 gallons per year) and a drop in the overall model fit (adjusted  $R^2 = 0.740$ ). However, most tellingly, the B values were off significantly from what one would expect (ex. Variable POOL B = 27.8; yearly evaporation in gallons per year is far in excess of this). Based on these findings it was decided that the APPROXINC variable should remain in the model.

## FESCUE

### Definition of Variable:

*Whether or not the turfgrass present at a residence is Fescue or an alternative turfgrass.* This is a binary (i.e., "dummy" in the vernacular) variable indicating presence (1) or absence (0) of a variable's specified condition.

### Results and Significance:

Fescue grasses (which are widely popular cool-season grasses found in local landscapes) have been observed to require large volumes of water in the Las Vegas area (ca. 91 inches), over 62% more annually than the other somewhat less popular warm-season Bermuda grass (requiring ca. 56 inches; calculations for both grasses are based on data from the local Cooperative Extension Office). Locally, Fescue is much less drought tolerant than Bermuda and has a correspondingly higher  $K_c$  value (the July  $K_c$  value for Fescue is calculated to be a very high 1.10 whilst only being ca. 0.71 for non-overseeded Bermuda; Source: University of Nevada Cooperative Extension Office).

Furthermore, being a cool-season grass, Fescue is capable of active photosynthesis all year long with sufficient irrigation and management, which is no doubt the reason for its desirability; it can yield an attractive green year round. Bermuda on the other hand usually goes into dormancy in the winter and it is likely many people curtail irrigation at dormancy so its total yearly application is even further reduced relative to Fescue. While there are of course different requirements for different types and morphologic forms of grasses (ex. tall vs. short fescue), the general finding that the cool-season grasses require more water than the warm season ones is well understood and this apparently translates into residences with Fescue having, on average, higher annual consumption at the household level ( $t=2.09$ ) (note: most residences had at least some turfgrass integral to their landscapes). Based on the multivariate analysis, a residence with Fescue may on average use more than 25,000 gallons more annually than one with a lower-water-use grass.

There is another possible inference that may be made. The submeter data is heavily dominated by Fescue landscapes and thus the highlighted gallons-per-square-foot application rates are probably at or near the actual *for Fescue*. It should be noted though that from the model, one might infer that in situations where there is not Fescue at the site, the B value of 59.1 may be the typical application rate, in gallons per square foot per year, for Bermuda installed at a residence. Though this derived value of 59.1 gallons per square foot per year (94.9 inches precipitation equivalents) is somewhat suppositional, and no doubt not exact given the standard error of the model, it appears to be a very reasonable average application rate that could be expected locally for Bermuda grass.

## PARCEL SIZE

Definition of Variable:

*The size, in square feet, of the parcels of study residences as specified in the Clark County Assessor's Office database.*

Results and Significance:

In the final version of the model, parcel size was technically not significant ( $t=1.79$ ); however, it was positively correlated with higher residential consumption in most multiple regressions developed so it is included here. It is reasonable to assume that, on average, residences associated with larger parcels are more likely to have higher consumption because they would be expected to have (i.) more, possibly lush, landscape (they are also more likely to have a pool) and (ii.) typically larger homes situated on them. Both of these would be anticipated to raise consumption due to larger residential landscapes having higher total outdoor irrigation requirements and larger houses being more likely to be inhabited by more or, perhaps, simply more heavily consuming, residents.

## POOL

Definition of Variable:

*The total water surface area of pools and spas in square feet at residences as measured by research personnel. For residences without pools this variable equates to zero.*

Results and Significance:

As with parcel size, pool surface area was not significant in the final most complete version of the model ( $t=1.70$ ), but often cropped up as significant in alternative models as being positively correlated with higher consumption. It is reasonable to include this variable as it is to be expected that the more evaporative water surface area outside at a residence owing to a pool and/or spa, the higher the evaporative water loss at the residence and the greater the need, in gallons, to replenish it.

## TOTALOCC

Definition of Variable:

*The total number of occupants at each study property in the analysis year (2001) as determined by survey.*

#### Results and Significance:

Though not a statistically significant independent variable in the final model ( $t=1.62$ ), and only occasionally significant in alternatives, the number of people living at the residences was ultimately included, as it lends explanatory strength to the model ( $\beta=0.524$ ) and it is logical to assume that consumption does increase with more people living at a location. That it is not statistically significant is actually a testament to the dominance of outdoor end uses in determining total yearly consumption at single-family properties in this region.

#### TOTALLAN

#### Definition of Variable:

*The total landscapeable area at a property.* This includes areas with landscape as well as areas potentially landscapeable.

#### Results and Significance:

This variable is difficult to interpret and was not significant in this particular model ( $t=-1.41$ ). The only reason for its inclusion is the sheer number of times it cropped up as significant in different alternative models. Here, however its sign is inverse of what would be anticipated (that greater landscapeable area would lead to higher consumption). It may be that it captures the inverse of the building and hardscape footprints, but this is only theory.  
check from here on...

#### **Variability in Annual Consumption for Irrigation of Monitored Xeric Landscape**

A model of yearly consumption for the monitored xeric component of landscapes for XS Group homes was also developed to attempt to evaluate the impacts of variables listed in the Scope (Appendix 1). The developed model has a much lesser fit than the total consumption model (adjusted  $R^2=0.40$ ), in part, one speculates, because other important but non-quantified or hidden variables are not included (one possible example – detailed data on controller management which may be more associated with management of turf rather than xeric areas). For this reason, no attempt is made to quantify impacts in a multivariate context as above, but rather the goal is to identify variables likely associated with xeric area consumption (for some attempts at quantification using univariate approaches consult Sovocool and Rosales<sup>11</sup> 2001).

Despite the limitations due to the weaker model, many variables did appear significant in most if not all modeling attempts, and these are discussed below in a format similar to the above discussion on annual consumption. The same strength of association denotation as used for the annual consumption model is applied to the xeric areas variable discussion as well. See introduction to *Sources of Significant Variability in Single-Family Residential Consumption* for more information.

## TOTALCAN

### Definition of Variable:

*The total canopy coverage in the monitored xeric area of the XS Group properties, in square feet.* This is calculated by first taking the observed plant diameters from the 2001 site review, dividing this number by two to get radius, then applying the formula for getting the area of a circle ( $A=\pi r^2$ ). This area result is then multiplied by the quantity of those plants observed to be at that size. The summation of all areas of all plants of all size classes in the study area is the total canopy coverage.

### Results and Significance:

It is reasonable to expect that total plant canopy coverage within the monitored xeric area would positively correlate to the total amount of water applied to that area as plant leaf surface area (evapotranspirational area) is the principal locale of water loss from vegetation. To replace this loss, areas with higher plant coverage should theoretically require more water and it should be expected that residents would respond by irrigating these more (via both longer run times and having irrigation systems of greater application capacitance). Examination for a link between total canopy coverage and total yearly consumption for xeric areas in a multivariate context confirms a significant association ( $t=4.31$ ; the relationship between coverage and per unit area consumption was also noted and explored in Sovocool and Rosales<sup>11</sup> 2001). One acknowledgement; this is a relatively simplistic finding, which does not fully explain the relationship between consumption and the taxa present and species' specific water use characteristics (this was beyond the practical scope of this investigation). Data on specific xeric species' water requirements is needed for this and this area remains worthy of more in-depth research.

## AVGFLOWR

### Definition of Variable:

*The arithmetic average flow rate, in gallons per minute, of all irrigation stations servicing monitored xeric landscape for each of the XS Group properties.*

### Results and Significance:

It has long been suspected that within the range of lower flow types of irrigation systems used to irrigate xeric areas, those capable of delivering water relatively faster via high-flow emitters may contribute to higher water consumption, especially when used by someone less knowledgeable about how to irrigate with different types of emitters. For this reason, SNWA's current Water Smart Landscapes program limits individual emitters to a maximum output of 20 gph as part of the program requirements (Appendix 5). Based on this research, this concern appears well-placed as the model shows stations with higher average flow rates are indeed associated with higher consumption in this study ( $t=4.14$ ). Typically, such station configurations may have one or more of the following conditions: sprays used for xeric-area irrigation, incorporation of high-flow emitters (such as turf bubblers), use of microsprays, stations composed of mixed types of irrigation emitters, and individual stations irrigating large and/or lush expansions of xeriscape (an exploration of how emitter class relates to average flow rates also appears in Sovocool and



Rosales<sup>11</sup> 2001; this manuscript suggested a strong association between irrigation system design and xeric area consumption as well).

## STUDYA

Definition of Variable:

*The xeric study land area (in square feet ) monitored via submeter for XS Group properties.*

Results and Significance:

It is logical to assume that, on average, the more area monitored by the submeter, the greater the consumption through that meter, and the significant association between monitored xeric-study area and total yearly consumption ( $t=3.08$ ) is consistent with this expectation (for further exploration of per-unit area savings, see *Comparison of Per-Unit Area Water Application between Turfgrass and Xeric Landscape*).

## TOTVAL

Definition of Variable:

*The dollar value of the residence as specified in the Clark County Assessor's Office database. This should not be considered the same as the home's market value.*

Results and Significance:

There was a positive association between the total value of the property and total consumption for xeric area consumption ( $t=2.94$ ). A discussion of how this variable tends to be positively associated with water consumption appears above in the discussion of the annual consumption model. It is worthwhile to again emphasize that given water use for residential landscapes can ultimately be considered discretionary, higher homeowners' wealth (here, evidenced by higher property value) may be anticipated to lead to greater consumption for landscape irrigation.

## PARCEL SIZE

Definition of Variable:

*The size, in square feet, of the parcel of a study residence as specified in the Clark County Assessor's Office database.*

Results and Significance:

The parcel size of the residence was significantly inversely associated with consumption for xeric area irrigation ( $t=-2.78$ ). This result was unexpected, as a relationship or mechanism acting to result in a link between parcel size and the irrigation of xeric areas on that parcel is not immediately obvious. The possibility that there is an inverse relationship between xeric study area and parcel area was examined, but this is not the case (rather, as would be expected, larger properties tended to be positively correlated with larger study areas, though this relationship is weak;  $R^2=0.064$ ). Likewise, the theory that perhaps larger parcels had xeric areas that might be sparser in terms of canopy was examined and rejected (the data does not support this).

Discussion and consideration of other findings led to some other possible explanations. One possibility is that those residences with larger parcels were more likely to incorporate native, lower-water-requirement plants in their landscapes. Some data supports the theory that owners of large properties may indeed make more use of native taxa, but only marginally so (the properties in the top 10% in parcel size had an average of 10.9% of their plant palette composed of native vegetation; the average for the rest of the properties was 6.9%).

Another theory is that larger xeriscape installations may be more likely to necessitate the need for a contractor, who is more likely to install a properly designed drip system and, as suggested by the findings linking station flow rate to consumption and (as revealed below) “drip-only” systems are more likely to result in lower total yearly consumption than those piecemealed together with multiple types of emitters. Since those residents doing larger xeriscape conversion projects were found to be more likely to use a contractor, there is some evidence supporting this second theory.

Perhaps the most likely reason for this finding is that people with smaller parcels are more able to afford to consume more water outdoors. To understand this reasoning better, consider an example of two sets of land, one acre each, in a similar area and climate each with all landscapeable area landscaped. One has a single residence upon it, the other acre, more subdivided, supports five homes (and thus is composed of five parcels). One would conclude, usually correctly, that the outdoor consumption for the *total area* would be greater for the one-home case, owing to its maintaining a greater amount of landscaped area (more of the available area is consumed as development in the five-homes case). But what about total water consumption for irrigation on a *per-parcel* basis? Each of the family income streams in the five-homes-per-acre case support less irrigated area than that for the single home on the one acre. Thus, each of these five owners can afford to support more discretionary water use as their respective landscape irrigation “shares” are less than for the one owner supporting all of that area. As a result, the owners of the smaller parcels may use more irrigation water per parcel than in the alternative case, and this may be what is being observed here (internal research by SNWA has shown that subdivision tends to result in higher per-parcel usage while decreasing usage for the total equivalent area).

Without more information, these are only hypotheses. At this time, while the inverse relationship between parcel area and xeric area consumption stands, the mechanism behind the relationship is not completely understood.

## DRIP

### Definition of Variable:

*Presence (1) or absence (0) of an exclusively drip irrigation system irrigating the xeric study area.* This is a binary variable.

### Results and Significance:

This is a different type of measure of the influence of irrigation system design on total xeric area water application. Specifically evaluated was whether the presence of a “true” drip system (no bubblers, microsprays, mixed systems) was associated with xeriscapes with lower consumption than others. The model does support this theory, with a significant finding that such “drip-only”

xeriscapes do have lower consumption ( $t=-2.27$ ). As suggested by Sovocool and Rosales<sup>11</sup> 2001, such systems typically have the lowest flow rates (average per-station flow rate = 4.0 gpm) of the types used to irrigate xeric landscape, so if run similar amounts of time to other systems, it would be expected that these would output lower total volume over a year. Based on the data, it does seem likely that many residents run their systems without careful consideration as to which kind of emitters they actually have, in turn resulting in systems composed exclusively of true drip emitters being associated with the least amount of water consumed over the year. Since slow application rates are generally the preference in irrigating drought-tolerant vegetation (this is especially the case with woody taxa) and because landscapes with “true” drip systems had the lowest consumption, this finding may be worthy of future considerations relevant to SNWA’s Water Smart Landscapes program.

### DONTKNOW

#### Definition of Variable:

*Whether or not the respondent was knowledgeable about the level of enforcement of local restrictions designed to reduce water waste.* This binary variable indicating presence (1) or absence (0) of understanding was adapted from part of an alternative answer to a question asking respondents if they felt enforcement of water waste provisions was “too lax,” “good,” or “too strict.” In addition to these responses, residents taking the survey were also given the option of answering “Don’t Know” if they did not have any sense of how aggressively water waste regulations in the area were enforced.

#### Results and Significance:

Theoretically a person’s viewpoints on water waste enforcement could tie into how diligently they practice good irrigation management. Recognizing this, the study staff formulated a question addressing this for the survey implemented in 2001. In preliminary analyses (Sovocool<sup>12</sup> 2002) there really was not a difference in per-unit area irrigation for xeriscapes between those respondents answering “too lax” and “good” (only two people said enforcement was “too strict” resulting in no ability to tie this to consumption with any statistical precision, though this is quite telling of how the community viewed enforcement in 2001). However, interestingly there was a difference between respondents with any kind of an opinion and respondents who had no sense of the issue. This suggested at the time that *awareness* of enforcement of water waste regulations may be a principal motivator to conserve, regardless of one’s viewpoint on how appropriate the level of enforcement is. The recurrence of this basic result, here in a multivariate scheme (i.e., those answering “don’t know” were associated with higher consumption in the regression model;  $t=2.13$ ) suggests that sensitizing the public about enforcement of water waste restrictions may be a powerful motivator for achieving outdoor water conservation.

### **FINANCIAL SAVINGS ASSOCIATED WITH CONVERSION PROJECTS AND COST EFFICIENCY**


As explained in the methods section, the research scope included a mandate to study some of the economics of xeriscape conversions, as this has been left relatively uninvestigated to date. Specifically, the directives were to quantify costs associated with the conversion and the subsequent maintenance of the xeriscape and to develop estimates as to what incentive level

would theoretically be necessary to entice people into doing conversion projects. Collection and analysis of this data is explained in *Methodology*, below, and in *Appendices 5* and *6*. Results are as follows below, starting with the conversion costs findings.

The average cost of the conversion for those converting in the XS Group was obtained via data collected on parts and materials, as well as contractor receipts. The average cost for all participants was \$2,881.21 for 1,862.1 sqft converted (\$1.55 per square foot for 91 participants sampled). The average cost for those who did the conversion themselves was \$2,428.31 for 1,766.22 sqft (\$1.37 per square foot), and the cost for those hiring a contractor was \$4,076.88 for 2,115.22 sqft (\$1.93 per square foot). These dollar amounts for costs and dollar valuations are as they stood in the late 1990s and have likely climbed slightly by today. As might be anticipated, it appears that residents may on average be more likely to hire a contractor for larger conversion projects.

Landscape maintenance requirements constitute a significant cost in labor and dollars directly spent. The relative amount of xeriscape at a residence figured prominently in landscape maintenance reductions for both these costs (Figure 12). For those who had at least 60% of their landscapeable area as xeric landscaping, maintenance savings of about one-third were realized versus those with 60% or more turf. The average difference is 2.2 hours/month in labor and \$206 per annum in direct expenditures (N=216). Landscape maintenance savings are value added on top of water bill savings. This serves to greatly enhance the attractiveness of xeriscape to the customer. Hessling<sup>12</sup> (2001) provides a detail of the capital costs and savings obtained.


**FIGURE 12: Average Monthly Maintenance Time and Annual Direct Expenditures for Participants Having At least 60% Turf or Xeriscape**



Bill savings for a typical mid-consumption range customer were modeled as explained in *Methodology* and in Appendix 4. Results show that there is a large difference in the monthly bills between a modeled residence with and without the conversion throughout the majority of the year (Figure 13). The total difference in the annual cost for water between these two homes using the current (2004) rate structure is \$239.92 - a significant savings attributable to the conversion (nearly \$0.15 per square foot converted per annum). It should be noted that this savings of 54% in total annual water charges is greater than would initially be anticipated from consumption savings data (Figure 6). This is because the Las Vegas Valley Water District, as well as the other SNWA member agencies, uses a tiered, increasing block rate structure.

Increasing block rate structures (also called conservation rate structures) are setup such that the more a user consumes on an average daily basis within a cycle, the more expensive, per unit (i.e., per gallon), water becomes. The high per-unit area application to turfgrass results in residences with more grass typically crossing thresholds into higher billing rate strata much more frequently and this in turn exacerbates their water costs per unit and, ultimately, their total costs. In this case, the difference in per-unit water charges for the two fifth-decile homes, with all charges considered over the entire year is about \$0.28 per thousand gallons (i.e., there is a 13% difference; effective prices of \$1.85 vs. \$2.13 per thousand gallons, respectively). The comparison highlights the utility of tiered rate structures as a conservation tool and for promotion of xeriscape as a conservation tactic.


**FIGURE 13: Modeled Monthly Water Bill for a Typical Las Vegas Area Home and The Same Home with an Average-Size Conversion**



The expected water bill savings a resident of a typical home would realize from doing an average-size conversion of turfgrass to xeriscape (anticipated monthly savings – including tier rate impacts) is thus as illustrated in Figure 14. As can be seen, the typical monthly water bill savings range from a low of \$5.68 (25%) in December to a high of \$40.84 (70%) in July, again reemphasizing that the greatest savings obtained by having xeric landscape are realized in the extremes of summer in this area. The savings obtainable serves to create a strong price signal to convert, especially when coupled with the incentive offered by SNWA currently (\$1.00 per square foot for qualifying residential conversions).

As suggested by Figures 13 and 14, on average xeriscape not only results in significant savings in water utility charges, it also makes the charges more manageable as they no longer “peak” to anywhere near the extent they did under the “no-conversion” condition. For the “no-conversion” model, the low-consumption month to high-consumption month ratio is 1:2.93 (the peak month is July). For the same house with the conversion, the ratio is 1:1.58 and the peak is pushed out to September owing to the difference in xeric irrigation pattern (Figure 8). For homes proximal to the modeled condition, xeriscape conversions appear to make paying monthly bills easier because the peak is (i.) greatly attenuated and (ii.) potentially pushed out until later in the year, so it does not parallel other local utility bills which peak in the summer (power, for example).

**FIGURE 14: Modeled Monthly Water Bill Savings for A Typical Las Vegas Area Home Completing an Average Size Conversion**



## ESTIMATED APPROPRIATE LEVEL OF FINANCIAL INCENTIVE

### Homeowner Perspective

Hessling<sup>13</sup> (2001) performed analyses of the financial viability of SNWA's xeriscape conversion program, "Southern Nevada Xeriscapes" (since revised and renamed to "Water Smart Landscapes"). It should be noted that at the time Hessling did his analysis, the program paid recipients an incentive of \$0.40 per square foot. He presented a Net Present Value (NPV) analysis demonstrating that, from the homeowner perspective, the return on investment by SNWA's conversion facilitation program is two to three years for a resident and that the incentive is not really required to induce change as the NPV is positive, even when no incentive is rewarded. See Hessling's manuscript for additional details.

A constructed model (Appendix 5) using a similar approach (and more recent data) seems to support the finding that no incentive is theoretically necessary for a typical do-it-yourself xeriscape conversion where subsequent financial savings in landscape maintenance are realized. However, the incentive may be important in a variety of other situations. The scenario, similar to the one used by Hessling as well as others, was explored by the model developed by SNWA (Appendix 5). Some of the most common scenarios explored, with findings from model outputs, are summarized in Figure 15.


In Figure 15, there are four different scenarios modeled (see explanation below), and each scenario has four associated results (*Methodology* and Appendix 5). The outputs associated with each exercise are: the average payback time (at a dollar per square foot) for a typical home doing a typical conversion (see Appendix 5), the average payback time without an incentive, the incentive required for a 3-year return on investment (ROI), and the incentive required for a 5-year ROI. Special note should be made regarding the expression of payback times. The display is not the range of payback times given the combination of scenario conditions, rather, it reflects that the expected average payback time falls sometime between the years shown. The model is based on annual, not monthly data thus the need to display outputs in this manner. The "incentive required" outputs, are simply average model outputs and are not to be considered appropriate for any one condition; their value is principally in comparative analyses between scenarios and in broad generalizations.

The summary (Figure 15) is designed to facilitate inferences about the economics of the conversion project. Along the horizontal axis are the "Only Conversion Material Costs" and "Conversion Material Costs + Labor" titles. The first scenario condition refers to situations where only the direct costs for materials, supplies, rentals, and other such items are considered. Residents doing their own xeriscape conversion might consider this to be their scenario if they consider only the real financial outlays paid and don't consider their time spent on the conversion to be a real financial cost. In contrast, the "Conversion Material Costs + Labor" condition includes a valuation of the time to actually do the conversion, which naturally lengthens the payback time. This perspective is more appropriate for those who consider the labor outputted by

themselves to be a true financial expenditure. It is also the appropriate model perspective to consider if the project is performed by a contractor.

Along the vertical axis of Figure 15, are the titles “Only Maintenance Goods Conserved” and “Conserved Maintenance Goods and Labor.” Similar to above, the “Only Maintenance Goods Conserved” condition reflects consideration of savings associated with only direct expenditures on things such as fertilizer, replacement irrigation parts, occasional replacement of capital items such as shovels, etc. (so long as the conversion is significant enough to yield savings in these areas; see the discussion surrounding Figure 12). The category “Only Maintenance Goods Conserved” would be most appropriate for people who do not consider the savings in labor obtained by having some of their area as xeriscape to be equivalent to a monetary outlay, situations where not enough of the total landscape area is converted to obtain this type of savings, or when a landscape maintenance company, which may or may not realize the savings, is either unwilling or unable to pass on labor savings to the customer as realized by the landscape retrofit. Again, there is an alternative category for the consideration of realized maintenance savings in labor costs resulting from the conversion. The maintenance savings plus labor savings category, “Conserved Maintenance Goods and Labor,” is most appropriate when enough of the yard has been converted that real savings in maintenance labor can be realized and when the owner attaches value to this. It would also be appropriate when the homeowner’s landscape company passes on realized labor savings to him or her.

The matrix of results developed (Figure 15) permits some inferences to be made about what scenarios turn around financially the fastest and are thus most readily facilitated by a landscape conversion incentive. In increasing order of time to payback (i.e., the first bulleted scenario is the most readily facilitated) these are:



Considering that the optimal price point for the first three of these scenarios is probably covered by the current incentive level, but not the old \$0.40-per-square-foot incentive, it may be that the SNWA hit upon a critical threshold value in stimulating the marketplace when it went to the



\$1.00 per-square-foot level in 2003. A recent surge in program interest in the residential sector is consistent with this (Appendix 5). Even in the fourth scenario, the current incentive level shortens the payback time such that the project might be deemed affordable by many people (see the associated 5-yr ROI). While few, if any, residents do a detailed economic assessment of the payback time for their respective xeriscape conversion projects, the dollar per square-foot is almost certainly perceived to make conversion projects much more “affordable,” plus there is significant symbolic value to the \$1.00-per-square-foot figure versus the past sub-dollar incentive levels.

If the payback time outputs presented in this model are close to reality, it may be that SNWA’s Water Smart Landscapes program will continue to experience high interest until a point where materials, supply (i.e., practically convertible turf), or services associated with the conversion project come to be in short supply and/or become expensive enough to cause feedback such that program enrollment is slowed.

**FIGURE 15: Summary of Economics of Typical Single-Family Xeriscape Conversion Projects Under Different Scenarios**

	<b>Only Conversion Material Costs</b>	<b>Conversion Material Costs + Labor</b>
<b>Only Maintenance Goods Conserved (or when labor savings not realizable)</b>	Avg. Payback Time at \$1.00 per SqFt: <b>3-4 Years</b>	Avg. Payback Time at \$1.00 per SqFt: <b>5-6 Years</b>
	Avg. Payback Time Without Incentive: <b>5-6 Years</b>	Avg. Payback Time Without Incentive: <b>8-9 Years</b>
	Incentive Required for 3-Year ROI: <b>\$1.03/SqFt</b>	Incentive Required for 3-Year ROI: <b>\$2.23/SqFt</b>
	Incentive Required for 5-Year ROI: <b>\$0.14/SqFt</b>	Incentive Required for 5-Year ROI: <b>\$1.34/SqFt</b>
<b>Conserved Maintenance Goods and Labor</b>	Avg. Payback Time at \$1.00 per SqFt: <b>1-2 Years</b>	Avg. Payback Time at \$1.00 per SqFt: <b>2-3 Years</b>
	Avg. Payback Time Without Incentive: <b>2-3 Years</b>	Avg. Payback Time Without Incentive: <b>4-5 Years</b>
	Incentive Required for 3-Year ROI: <b>None Req.</b>	Incentive Required for 3-Year ROI: <b>\$0.91/Sqft</b>
	Incentive Required for 5-Year ROI: <b>None Req.</b>	Incentive Required for 5-Year ROI: <b>None Req.</b>

## SNWA Perspective

The financial viability of SNWA's Water Smart Landscapes Program is difficult to assess as resource alternatives available to the Authority against which this "water option" may be measured are diverse and have widely divergent respective costs (SNWA<sup>20</sup> 2003). Furthermore, availability of water resources is not constant and shortage or surplus conditions can exist which can make using these as standards against which conservation programs can be measured again difficult. A prime and current example of this is the drought that the Lower Colorado River Basin is experiencing which is currently impacting SNWA (SNWA Drought Plan<sup>21</sup> 2003). In these types of situations, the economics of conservation programs are obviously enhanced, and it is against this backdrop that the economics of the Water Smart Landscapes Program is being considered in this study.

In Hessling's analyses<sup>13</sup>, the drought had not yet been recognized and designated as such and SNWA had no drought policies in place at the time of the analysis. He grounded his analysis in comparing the marginal cost of water in the Southwest to the marginal benefit realized by the incentive program. In doing so, he concluded that the cost of the incentive program at the time was just offset by its resource value, and the program was thus a worthwhile initiative (see analysis for details).

In 2004, a reanalysis of the Water Smart Landscapes Program was done to consider the economic viability of it in the face of the drought and the current resource and program incentive values. Given the current scarcity of local water resources, the drought, and the fact that SNWA is now approaching the point of withdrawing its full Colorado River allotment (SNWA<sup>20</sup> 2003), the Las Vegas Valley Water District has recently paid \$9,500 per acre-foot for undeveloped groundwater rights in the local basin and, furthermore, views this purchase as a bargain (LVVWD<sup>22</sup> 2003). Because the largest purveyor member in the SNWA is willing to pay this amount currently for undeveloped, non-administered water rights, this should be a good alternative price for comparing the cost effectiveness of the program on a per-square-foot basis (not including administrative and advertising costs).

It follows that to estimate the savings yielded by the program in dollars per square foot, the above marginal cost of water, converted to a square-foot basis, can be multiplied by the savings per square foot yielded by the conversion as below:

$$\text{\$9,500 per acre-foot} \times 325851 \text{ gallons per acre-foot} \times 55.8 \text{ gallons per sqft yield} = \text{\$1.627 per sqft}$$

The cost calculation is slightly more complex, as the SNWA not only spends the \$1.00 per square foot to incentivize the conversion, but it also forgoes the yield it would have claimed on this amount had it invested it. The mature yield of municipal bonds in February 2004 is used as this alternative rate. Thus the true cost per square foot for SNWA can be estimated as:

$$\text{\$1.00 per sqft expended} + (\text{\$1.00} + 4.65\% \text{ mature interest yield if invested instead}) = \text{\$1.047}$$

The cost-effectiveness of the program can then be calculated as the difference between these values:

$\$1.627 \text{ per sqft saved} - \$1.047 \text{ per sqft saved} = \$0.58 \text{ per sqft net positive value to SNWA}$

The analysis suggests that for each dollar the SNWA is spending for the incentive, it is bringing in \$1.58 and that the program appears as such to be a good deal from a financial perspective for SNWA. The ca. 37% net positive value means the program should be financially advantageous even with addition of the program advertising and administration costs which have not to date been quantified.

In considering the theoretical maximum that SNWA could pay for the program (a component of Objective 6), it should be noted that \$1.627 is not the maximum as, again, the yield of the alternative investment must be considered. Subtracting out this missed or forgone yield results in a figure of \$1.55 and this is the theoretical maximum price SNWA could currently justifiably sustain. Again, the actual maximum would be anticipated to be lower due to program administration costs.

## Executive Summary and Conclusions

The major conclusions of this research are as follows:

1. Xeriscape conversion projects can save vast quantities of water at single-family residences. Homes in this study saved an average of 96,000 gallons annually following completion of an average-size conversion project. This is a savings of 30% in total annual consumption; a finding in line with those yielded by other research studies in this region.
2. Over the long timeframe of this study, total yearly savings have neither eroded nor improved across the years. On average, household consumption drops immediately and quickly stabilizes.
3. There is an enormous difference in application of water to locally used turfgrasses and xeric landscape by residents. On average, each year residents applied 73.0 gallons per square foot (117.2 inches) of water to grow turfgrass in this area and just 17.2 gallons per square foot (27.6 inches) to xeric landscape areas. The difference between these two figures, 55.8 gallons per square foot (89.6 inches) is the theoretical average savings yielded annually by having xeriscape in lieu of turf in this area. This is a *substantial* savings (76.4%) when considered in the context of the available residential water conservation measures. A sub-study of other commercial properties with xeriscape found the average application to xeric areas by these customers to be essentially equivalent to that observed for the residential customers.
4. Over the course of a year, the difference in application between turf and xeric areas varies in a predictable bell-shaped-curve manner, with the greatest difference occurring in summer. This is because turf irrigation peaks to a much greater extent in summer than xeric irrigation. The difference in irrigation between these two types of landscape varies from as little as 1.56 gallons per square foot for the month of December, on up to 9.62 gallons per square foot for the month of July.
5. In comparing irrigation application to the reference evapotranspirational rate ( $ET_0$ ), it was found that on average application to turf exceeded  $ET_0$  in every month except March, exceeding it the most May through November. In contrast, xeric application remained well below  $ET_0$  year round.
6. The author experimented with using a locally invoked “rule-of-thumb” which holds that xeric plantings require about a third of the evapotranspirational rate as needed for turf. In comparing this developed reference,  $0.33(ET_0)$ , to application, it was found that these values were, in absolute terms, somewhat close month to month and very close over the entire year. In comparing this developed reference to application, it was found that xeric application was below  $0.33(ET_0)$  half the year and above it the other half of the year (September-February).

7. Relative to questions about irrigation management and the potential for further efficiency gains, findings associated with conclusions 4 through 6 and subsequent analyses led the author to suggest that (i.) the greatest absolute savings from assorted improvements in irrigation will be realized in the summer, but (ii.) the most readily obtained efficiency improvements (i.e., not requiring capital outlays) yielded from better controller management may be obtained September through January, as this is the period when a lot of residents fail to successfully decrease irrigation in response to lower irrigation requirements (for both types of landscape).
8. Multivariate regression modeling was used to help discover some of the factors associated with variability in water consumption at single-family residences. These are:
  - i. The amount of turf at the residence (positive correlation).
  - ii. The property value of the residence (as indicated by the local assessor's office; positive correlation).
  - iii. The age of the residence (positive correlation).
  - iv. The total income of the property's residents (positive correlation).
  - v. Whether or not the turfgrass present at the residence is Fescue (a locally popular cool-season grass; positive correlation). As a side-result from one of the multivariate analyses, Bermuda grass may be receiving approximately 59 gallons per square foot per year – certainly less than the application for the much more common cool-season grass in this study.

Some variables which were significant in many other incarnations of the model (but not the final model) include parcel size, surface area of open water for pools and spas, the total number of occupants living at the residence, and total landscapeable area.

9. A similar approach was used to identify some of the factors associated with variability in irrigation application to monitored xeric areas. These are:
  - i. The total canopy coverage within the xeric area (positive correlation).
  - ii. The average per-station flow rate of the installed irrigation system serving the xeric area (positive correlation).
  - iii. The size of the xeric area (positive correlation).
  - iv. The property value of the residence (positive correlation).
  - v. Parcel size (inverse correlation).
  - vi. Whether or not the irrigation system was exclusively a drip irrigation system (i.e., not composed of microsprays, bubblers, other higher flow emitters, or combinations of emitters; inverse correlation).
  - vii. Whether or not the resident responsible for managing irrigation at the home is knowledgeable about enforcement of local provisions prohibiting outdoor water waste (inverse correlation).
10. Tracking of the costs residents incurred when converting their landscapes from turf to xeric landscape revealed that at the time of the study, the average conversion cost was \$1.55 per square foot across all of the conversion projects for which data was available. The average cost for those who did the work themselves was \$1.37 per square foot, and for those employing a contractor, it was \$1.93 per square foot. All of these costs are probably higher today due to inflation and a strong market for conversion projects.

11. In comparing those with 60% or more of their landscape as xeric landscaping and those whose landscape was 60% or more turf, it was found that those with the majority as xeriscape condition enjoyed a 2.2 hrs-per-month reduction in landscape maintenance and an additional \$206 per annum savings in direct maintenance expenditures as well. This represented a savings of about a third in total landscape labor and maintenance expenditures, respectively.
12. A model of two identical homes, one near the average for consumption (technically in the fifth decile for consumption), the other the same, but having completed an average-size conversion, revealed the following:
  - i. The annual water bill savings yielded by landscape conversion projects can be large. For the Las Vegas Valley Water District customer modeled, the annual financial savings was \$239.92 (figure includes all applicable surcharges). This equates to a savings of nearly \$0.15 per square foot.
  - ii. This is a large savings of 54% in total annual charges for water consumption. This level of savings is elevated over what might have been initially anticipated due to an aggressive tiered water rate structure. The effective average fifth-decile annual water charges with all surcharges added would be \$2.13/kgal for the typical traditional home and \$1.85/kgal for the one having completed the average-size conversion.
  - iii. The savings vary by season as expected by the findings associated with the submeter data. Whereas the bill payer of the home having done the conversion saved 25% (\$5.68) in charges for December vs. the typical homeowner, the same individual would realize an enormous savings of 70% (\$40.84) for July. One of the great benefits of xeriscape is that it drastically mediates “peaking” in summer, making summer bills much more affordable for households, especially since power bills also peak in summer.
13. A model was also created to explore payback time and the appropriateness of the financial incentive. This revealed that payback time varies in part on whether or not homeowners do the work themselves or enlist the assistance of a contractor and whether or not savings in maintenance labor is actually realized. Modeling proceeded such that different combinations of these scenarios were explored. The results suggest that in most situations the current SNWA incentive is sufficient to help facilitate conversions such that there is an acceptable time to return on investment (ROI) for the homeowner. In order of increasing time to ROI from the point of conversion, with a dollar-per-square foot incentive from SNWA, these are as follows:
  - Situations where only the material costs of the conversion are valued and where savings in both maintenance goods and labor can be realized (average payback time of one to two years).
  - Situations where both the material costs of the conversion and the labor cost of the conversion are valued and where savings in both maintenance goods and labor can be realized (average payback time of two to three years).

- Situations where only the material costs of the conversion are valued and where only savings in maintenance goods (not labor) can be realized (average payback time of three to four years).
- Situations where both the material costs of the conversion and the labor cost of the conversion are valued and where only savings in maintenance goods (not labor) can be realized (average payback time of five to six years).

14. An economic analysis of the cost-efficiency of SNWA's Water Smart Landscapes Program suggests that in theory the program is cost-efficient and could be bringing in the equivalent of \$1.58 for each \$1.00 spent on rebate incentives (a 37% positive return) by way of effectively freeing up local water resources for immediate use. When the opportunity cost is included in the calculation, it is determined that the theoretical maximum incentive SNWA should be currently willing to pay in 2004 dollars is \$1.55 per square foot (the actual maximum is less due to program administration costs). In practice, this means it is probably not cost-effective to raise the incentive further at this time as the level necessary to yield a 3-yr ROI for those not yet facilitated to convert (i.e., the final bulleted scenario in Conclusion 13) equates to \$2.23, an incentive level far in excess of the theoretical top-out point for an incentive provided by SNWA. Furthermore, in the majority of modeled scenarios, the dollar per-square-foot is sufficient incentive for homeowners to justify the landscape conversion project.



## References

1. Bent, 1992. East Bay Municipal Utility District. CA (data recapitulated from citation in Reference 4).
2. Testa, A. and Newton, A., 1993. An Evaluation of a Landscape Rebate Program. *AWWA Conserv'93 Proceedings*, December. 1763 – 1775. Mesa, AZ.
3. Nelson, J., 1994. Water Saved By Single Family Xeriscapes. *1994 AWWA Annual Conference Proceedings*, June. 335-347. North Marin Water District, Novato, CA
4. Gregg, T. et al., 1994. *Xeriscaping: Promises and Pitfalls*. City of Austin. Austin, TX.
5. Colorado River Compact of 1922. United States of America. H. Doc 605, 67<sup>th</sup> Cong., 4<sup>th</sup> Sess.
6. Boulder Canyon Project Act. 1928. United States of America. H. Doc 642, 70<sup>th</sup> Cong., 2<sup>nd</sup> Sess.
7. Cooperative Agreement No. 5-FC-30-00440, 1995. Cooperative Agreement Between Bureau of Reclamation and Southern Nevada Water Authority for Research Investigation of Implementation of Xeriscape.
8. Mayer, P. and DeOreo, W. et al., 1999. Residential End Uses of Water. AWWA Research Foundation. U.S.A.
9. DeOreo, W., Mayer, P., Rosales, J., 2000. Xeriscape Conversion for Urban Water Conservation. *National Irrigation Symposium Proceedings*, November. Phoenix, NM.
10. Sovocool, K., Hessling, M., and Rosales, J., 2000. Residential Xeriscape: Answering Some Common Questions on Water and Landscape Maintenance Savings by Field Research. *New Mexico Xeriscape Conference 2000 Proceedings*, October. Albuquerque, NM.
11. Sovocool, K. and Rosales, J., 2001. A Five-year Investigation into the Potential Water and Monetary Savings of Residential Xeriscape in the Mojave Desert. *2001 AWWA Annual Conference Proceedings*, June. Washington D.C. Southern Nevada Water Authority, NV.
12. Sovocool, K., 2002. Subtle Mediators of Water Savings in Xeriscape Conversion Projects. *AWWA 2002 Water Sources Conference*, January. Las Vegas, NV. Southern Nevada Water Authority, NV.
13. Hessling, M., 2001. Turf Landscapes versus Xeriscapes: Analysis of residential landscapes in the Las Vegas Valley, Nevada. Master's Project. Duke University, NC.

14. Mayer, P., Towler, E., and DeOreo, W., et al., 2004. National Multiple Family Submetering and Allocation Billing Study. Aquacraft, Inc., and East Bay Municipal Utility District, CA.
15. Rosales, J., Weiss, C., and DeOreo, W., 2002. The Impacts of Submetering on Water Usage at Two Mobile Home Communities in Las Vegas, Nevada. *AWWA 2002 Water Sources Conference*, January. Las Vegas, NV. Southern Nevada Water Authority, NV.
16. *Predicting and Estimating Landscape Water Use*. The Irrigation Association. April 2001.
17. *A Guide to Estimating Irrigation Water Needs of Landscape Plantings in California – The Landscape Coefficient Method and WUCOLS III*. University of California Cooperative Extension and California Department of Water Resources. August 2000.
18. Kozlowski, T., Kraper, P., and Pallardy, S., 1990. *The Physiological Ecology of Woody Plants*. Academic Press, Inc., San Diego.
19. Gregg, T. et al., 1999. *Xeriscaping: Sowing the Seeds for Reducing Water Consumption*. City of Austin. Austin, TX.
20. *SNWA 2003 Water Resources Plan*. Southern Nevada Water Authority. 2003.
21. *SNWA Drought Plan*. Supplement to *SNWA 2003 Water Resources Plan*. Southern Nevada Water Authority. 2003.
22. *Agreement Between Ruffin Gaming, LLC, Ruffin Vegas, LLC and the District for the Purchase of 300 Acre-feet of Certificated and Permitted Groundwater Rights for An Amount Not To Exceed \$2,850,000*. Las Vegas Valley Water District Board of Directors Agenda Item #8. Las Vegas Valley Water District. April 2003.

## Appendices

**APPENDIX 1: ATTACHMENT A (SCOPE OF WORK) FOR BOR  
AGREEMENT 5-FC-30-00440 AS REVISED 11/19/98**

**APPENDIX 2: MULTIVARIATE MODEL DETAILS**

Note: Detailed definitions of variables and units for with each variable for both of the below models appear in the corresponding sections within *Sources of Significant Variability in Single-Family Residential Consumption*.

**TABLE 19: Multivariate Regression Model of Annual Single-Family Residential Consumption**

Regression Summary

Dependent Variable: MAINMETE (i.e., annual consumption registered through mainmeter)

$R^2=0.80889235$ ; Adjusted  $R^2=0.80046113$

$F(9,204) = 95.940$ ;  $p<0.0001$

Std. Error of Estimate=76890

Variable	Beta	Std. Error of Beta	B	Std. Error of B	t(204)	p - level
Intercept			-90852.6	25413.77	-3.57494	0.000437
POOL	0.060698	0.035627	51.3	30.13	1.70371	0.089959
TOTALTUR	0.622464	0.041887	59.1	3.98	14.86045	0.000000
TOTALLAN	-0.145252	0.102765	-5.5	3.90	-1.41344	0.159051
APPROXINC	0.073217	0.033839	0.3	0.14	2.16370	0.031649
FESCUE	0.068672	0.032854	25756	12322.71	2.09020	0.037839
TOTVAL	0.281661	0.051686	2.1	0.39	5.44950	0.000000
PARCELSI	0.214206	0.119536	5.9	3.28	1.79197	0.074620
NLTHOMEA	0.117091	0.043809	1600.6	598.85	2.67274	0.008132
TOTALOCC	0.52416	0.032356	8860.4	5469.42	1.61999	0.106780

**TABLE 20: Multivariate Regression Model of Annual Xeric Study Area Consumption**

Regression Summary

Dependent Variable: SUBMETER (i.e., annual consumption registered through submeter)

$R^2=.64787230$ ; Adjusted  $R^2=.41973852$

$F(7,178) = 18.394$ ;  $p<0.0001$

Std. Error of Estimate=32272

Variable	Beta	Std. Error of Beta	B	Std. Error of B	t(178)	p - level
Intercept			-7697.6	8973.436	-0.85782	0.392144
STUDYA	0.211132	0.068633	6.4	2.087	3.07623	0.002427
TOTALCAN	0.299352	0.069467	9.2	2.126	4.30934	0.000027
DONTKNOW	0.122082	0.57381	10922.2	5133.663	2.12756	0.034750
TOTVAL	0.213746	0.072592	0.4	0.137	2.94447	0.003667
PARCELSI	-0.211758	0.076239	-1.5	0.524	-2.77756	0.006064
AVGFLOWR	0.265679	0.064116	3637.4	877.802	4.14372	0.000053
DRIP	-0.133730	0.058997	-13615	6006.406	-2.26674	0.024609

### APPENDIX 3: RAW DATA

Raw data for possible further analysis is included in the file “BORdata.mdb.” A copy of this Microsoft Access database file is being included on disk with submission of this report to BOR. Below is the data description and dictionary for the file (this is also saved on disk).

#### Xeriscape Conversion Study Data Description

1. **tblCustomerList – 716 Records**, basic customer information.
  - a. ClientID – SNWA Customer identification number
    - i. Number – Long Integer
    - ii. Primary Key
  - b. Program – Indicates if the property is a xeriscape or turf study site
    - i. Text – 50
    - ii. XS = Xeriscape Study, TS = Turf Study
  - c. FirstName – Property occupant’s first name
    - i. Text – 50
  - d. LastName – Property occupant’s last name
    - i. Text – 50
  - e. Address – Address of property
    - i. Text – 50
  - f. City
    - i. Text – 50
  - g. Zip – Postal code
    - i. Text – 5
  - h. HomePhone
    - i. Text – 50
  - i. WorkPhone
    - i. Text – 50
  - j. Comments – Optional comment field
    - i. Memo
  - k. OwnerChange – Indicates if there has been a change in the ownership of the property.
    - i. Boolean
  - l. FollowupMonth – Number of the month the property has been assigned to schedule an annual follow-up site visit.
    - i. Text – 2
  - m. AccountNum – LVVWD / SNWA account number assigned to the property
    - i. Number – Long Integer
  - n. ServiceArea – Indicates if the customer receives service from LVVWD or one of the other entities.
    - i. Text – 50
    - ii. S = LVVWD Service, O = Outside Entity.

- o. Agreement – Date the customer signed the agreement to become a participant in the study.
  - i. Date/Time
- p. FinalReview – Date final inspection site visit was conducted after the installation of the submeter.
  - i. Date/Time
- q. Status – File quality status indication.
  - i. Text – 50
- r. FileQuality – Quality rating of file information
  - i. Text – 50

**2. tblAllSubmeterData – 2667 Records, customer’s submetered consumption data.**

- a. nltClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. nitYear
  - i. Number – Integer
  - ii. Primary Key
- c. txtEntity – Indicates which water provider services the customer
  - i. Text – 5
- d. txtProgram – Indicates if the property is a xeriscape or turf study site
  - i. Text – 2
  - ii. XS = Xeriscape Study, TS = Turf Study
- e. nstJan – January submeter consumption in gallons
  - i. Number – Single Precision
- f. nstFeb – February submeter consumption in gallons
  - i. Number – Single Precision
- g. nstMar – March submeter consumption in gallons
  - i. Number – Single Precision
- h. nstApr – April submeter consumption in gallons
  - i. Number – Single Precision
- i. nstMay – May submeter consumption in gallons
  - i. Number – Single Precision
- j. nstJun – June submeter consumption in gallons
  - i. Number – Single Precision
- k. nstJul – July submeter consumption in gallons
  - i. Number – Single Precision
- l. nstAug – August submeter consumption in gallons
  - i. Number – Single Precision
- m. nstSep – September submeter consumption in gallons
  - i. Number – Single Precision
- n. nstOct – October submeter consumption in gallons
  - i. Number – Single Precision
- o. nstNov – November submeter consumption in gallons
  - i. Number – Single Precision

- p. nstDec – December submeter consumption in gallons
  - i. Number – Single Precision
- q. nstTotal – Total yearly submeter consumption in gallons
  - i. Number – Single Precision

**3. tblAOX2 – 702 Records**, parcel information from Assessor’s database

- a. ClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. PLDECKSQF – Pool decking square footage
  - i. Number – Single Precision
- c. STORAGESQF – Storage area square footage
  - i. Number – Single Precision
- d. PAVE1SQF – Paved area one square footage
  - i. Number – Single Precision
- e. PAVE2SQF – Paved area two square footage
  - i. Number – Single Precision
- f. PATIO1SQF – Patio one square footage.
  - i. Number – Single Precision
- g. PATIO2SQF – Patio two square footage
  - i. Number – Single Precision
- h. PATIO3SQF – Patio three square footage
  - i. Number – Single Precision
- i. GARAGE1SQF – Garage area 1 square footage
  - i. Number – Single Precision
- j. GARAGE2SQF – Garage area 2 square footage
  - i. Number – Single Precision
- k. CARPORTSQF – Carport area square footage
  - i. Number – Single Precision
- l. FIRSTFLSQF – First floor footprint square footage
  - i. Number – Single Precision
- m. TOTALHS – Total of all hardscape areas
  - i. Number – Single Precision
- n. PARCEL – Assessor’s parcel number
  - i. Text – 11

**4. tblETDatawithCustomerIDs – 716 Records**, total monthly and annual evapotranspiration rates for 2001 by month correlated with SNWA client identification numbers.

- a. ClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. ETType
  - i. Text - 50
- c. JanET
  - i. Number – Single Precision



- d. FebET
  - i. Number – Single Precision
- e. MarET
  - i. Number – Single Precision
- f. AprET
  - i. Number – Single Precision
- g. MayET
  - i. Number – Single Precision
- h. JunET
  - i. Number – Single Precision
- i. JulET
  - i. Number – Single Precision
- j. AugET
  - i. Number – Single Precision
- k. SepET
  - i. Number – Single Precision
- l. OctET
  - i. Number – Single Precision
- m. NovET
  - i. Number – Single Precision
- n. DecET
  - i. Number – Single Precision
- o. TotalET
  - i. Number – Single Precision

**5. tblETDatawithCustomerIDsAvg – 716 Records**, average monthly and annual evapotranspiration rates for 2001 by month correlated with SNWA client identification numbers.

- a. ClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. ETType
  - i. Text – 50
- c. JanAvgET
  - i. Number – Single Precision
- d. FebAvgET
  - i. Number – Single Precision
- e. MarAvgET
  - i. Number – Single Precision
- f. AprAvgET
  - i. Number – Single Precision
- g. MayAvgET
  - i. Number – Single Precision
- h. JunAvgET
  - i. Number – Single Precision

- i. JulAvgET
    - i. Number – Single Precision
  - j. AugAvgET
    - i. Number – Single Precision
  - k. SepAvgET
    - i. Number – Single Precision
  - l. OctAvgET
    - i. Number – Single Precision
  - m. NovAvgET
    - i. Number – Single Precision
  - n. DecAvgET
    - i. Number – Single Precision
  - o. TotalAvgET
    - i. Number – Single Precision
- 6. tblInstalledCanopy – 447 Records**, total of square feet of plant coverage of xeriscape participants upon installation of the landscape.
- a. ClientID – SNWA Customer identification number
    - i. Number – Long Integer
    - ii. Primary Key
  - b. InstCanopyArea – Installed plant canopy square feet.
    - i. Number – Single Precision
- 7. tblParcelInfo – 702 Records**, Information from Clark County Assessor’s office database extracted November 2002.
- a. ClientID – SNWA Customer identification number
    - i. Number – Long Integer
    - ii. Primary Key
  - b. ParcelNum – Assessor’s office parcel number
    - i. Text – 11
  - c. ParcelSize – Size of parcel in square feet
    - i. Number – Single Precision
  - d. CONSTYR – Construction year
    - i. Number – Integer
  - SALEPRICE – Last Sales price
    - ii. Number – Long Integer
  - e. LYTOTAL – Last years assessed value land and improvement
    - i. Number – Long Integer
  - f. SALEDATE – Last sales date (Year)
    - i. Text - 6
  - g. nltHomeAge – Age of home calculated by construction year from the year 2001.
    - i. Number – Long Integer

- 8. tblResults – 603 Records**, collection of landscape areas, yearly water consumption data, other site, and customer information
- a. nltClientID – SNWA Customer identification number
    - i. Number – Long Integer
    - ii. Primary Key
  - b. Program – (TS = Turf Study Participant, XS = Xeriscape Study)
    - i. Text – 50
  - c. Converted – Area converted if XS participant
    - i. Number – Single Precision
  - d. Pool – Square footage of pool surface if present
    - i. Number – Single Precision
  - e. GardenMon – Square footage of garden area where the irrigation is monitored by the submeter
    - i. Number – Single Precision
  - f. GardenUnmon – Square footage of garden area where the irrigation is not monitored by the submeter
    - i. Number – Single Precision
  - g. Other – Square footage of other undeveloped property area. No irrigation, plants, or hardscape present.
    - i. Number – Single Precision
  - h. Study – Total xeriscape area where irrigation is monitored by the submeter. Applies to XS participant only.
    - i. Number – Single Precision
  - i. TurfMon – Square footage of turf grass where irrigation is monitored by the submeter.
    - i. Number – Single Precision
  - j. TurfUnmon – Square footage of turf area where the irrigation is not monitored by the submeter
    - i. Number – Single Precision
  - k. XeriMon – Square footage of xeriscape where irrigation is monitored by the submeter. (Applies to Turf Study Group)
    - i. Number – Single Precision
  - l. XeriUnmon – Square footage of xeriscape area where the irrigation is not monitored by the submeter.
    - i. Number – Single Precision
  - m. TotalLandscape – Total of all landscapable area on the property.
    - i. Number – Single Precision
  - n. TotalEvaporative – Total of all landscapable area with pool area added.
    - i. Number – Single Precision
  - o. dtt2001SR – Date of final annual visit conducted in 2001.
    - i. Date/Time
  - p. AgeOfXeriscape – Age of xeriscape in days calculated by the difference in days between the post submeter installation inspection and the final 2001 follow-up site visit.
    - i. Number – Long Integer

- q. TotalXeriArea – Total of all xeriscape areas, monitored and unmonitored.
  - i. Number – Single Precision
- r. Status – File quality status indication.
  - i. Text - 50
- s. TotalCanopy – Total of all plant canopy areas as of the 2001 annual site visit.
  - i. Number – Single Precision
- t. nitYear
  - i. Number – Integer
- u. txtEntity – Water agency that services the customer.
  - i. Text - 5
- v. Submeter2001 – Total gallons used in the year 2001 through the submeter
  - i. Number – Single Precision
- w. Mainmeter2001 – Total gallons used in the year 2001 through the main meter
  - i. Number – Single Precision
- x. pctGarden – Percent of total landscape area in garden
  - i. Number – Single Precision
- y. pctXeri – Percent of total landscape in xeriscape
  - i. Number – Single Precision
- z. pctTurf – Percent of total landscape area in turf
  - i. Number – Single Precision
- aa. pctOther – Percent of total landscape in other non-landscaped area
  - i. Number – Single Precision
- bb. pctPool – Percent of total landscape area in pool
  - i. Number – Single Precision
- cc. pctXeriRank – Xeriscape study participants were divided into ten percent ranges based upon percentage of landscape in xeriscape and given a ranking.
  - i. Number – Single Precision
- dd. XeriDensity – Percent of plant coverage per square foot of xeriscape.
  - i. Number – Single Precision
- ee. TurfType – Type of turf (Bermuda, Fescue, etc.) on property if present.
  - i. Text – 50
- ff. BarrierType – Type of weed barrier present if Xeriscape study participant.
  - i. Text – 50

- 9. tblSurveyInfoOfInterest – 603 Records**, Responses to survey questions. Each possible response is listed as a separate field. The responses are grouped where appropriate.
- a. CLIENTID – SNWA Customer identification number
    - i. Number – Long Integer
    - ii. Primary Key
  - b. SurveyAnswered – “Yes” or “No” Indicates if the customer answered any of the questions on the survey.
    - i. Text – 3
  - c. CLOCKADJ – How many times per year the irrigation clock was adjusted
    - i. Number – Byte

- d. INCBILL – How much of an increase in the monthly bill would produce conservation?
  - i. Number – Integer
- e. RESPAGE – Respondent’s age
  - i. Number – Byte
- f. Respondent’s gender
  - i. MALE
    - 1. Number – Byte (1 = Yes, 0 = No)
  - ii. FEMALE
    - 1. Number – Byte (1 = Yes, 0 = No)
- g. Respondent’s marital status
  - i. MARRIED
    - 1. Number – Byte (1 = Yes, 0 = No)
  - ii. SINGLE
    - 1. Number – Byte (1 = Yes, 0 = No)
  - iii. WIDOWED
    - 1. Number – Byte (1 = Yes, 0 = No)
- h. RETIRED – Indicates if respondent is retired or not
  - i. Number – Byte (1 = Yes, 0 = No)
- i. NATIVE – Native to southern Nevada?
  - i. Number – Byte (1 = Yes, 0 = No)
- j. AGE65PLS – Number of residents at the property age 65 and older
  - i. Number – Byte
- k. APROXINC – Median of a range of household income
  - i. Number – Long Integer
- l. Respondent’s opinion on Water Waste enforcement
  - i. DONTKNOW
    - 1. Number – Byte (1 = Yes, 0 = No)
  - ii. GOOD
    - 1. Number – Byte (1 = Yes, 0 = No)
  - iii. LAX
    - 1. Number – Byte (1 = Yes, 0 = No)
  - iv. STRICT
    - 1. Number – Byte (1 = Yes, 0 = No)
- m. Highest Education Level
  - i. ASSOCDEG – Associate’s degree
    - 1. Number – Byte (1 = Yes, 0 = No)
  - ii. BACHDEG – Bachelor’s degree
    - 1. Number – Byte (1 = Yes, 0 = No)
  - iii. GRADDEG – Graduate degree
    - 1. Number – Byte (1 = Yes, 0 = No)
  - iv. HSDEG – High school degree
    - 1. Number – Byte (1 = Yes, 0 = No)

- v. SOMECOLL – Some College
  - 1. Number – Byte (1 = Yes, 0 = No)
- vi. SOMEGRAD – Some graduate education
  - 1. Number – Byte (1 = Yes, 0 = No)
- vii. TECHTRAD – Technical or trade school
  - 1. Number – Byte (1 = Yes, 0 = No)
- viii. ADTECTRN – Advanced technical training
  - 1. Number – Byte (1 = Yes, 0 = No)
- n. Type of Grass at residence
  - i. BERMUDA
    - 1. Number – Byte (1 = Yes, 0 = No)
  - ii. FESCUE
    - 1. Number – Byte (1 = Yes, 0 = No)
  - iii. BUFFALO
    - 1. Number – Byte (1 = Yes, 0 = No)
  - iv. BFMIX – Bermuda / Fescue Mix
    - 1. Number – Byte (1 = Yes, 0 = No)
  - v. UNKNOWN
    - 1. Number – Byte (1 = Yes, 0 = No)
  - vi. NONE
    - 1. Number – Byte (1 = Yes, 0 = No)

**10. tblSurveyTotBath – 623 Records**, total number of bathrooms on the property

- a. ClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. Bathrooms
  - i. Number – Single Precision

**11. tblSurveyTotOccupants- 341 Records**, total number of occupants in the household at the time of the survey.

- a. nltClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. TotalOccupants
  - i. Number – Integer

**12. tblIrrigationData – 355 Records**, Irrigation system components for each property were assessed, and each property assigned to one of the following categories.

- a. ClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. AvgFlowRate – Average flow rate of all stations
  - i. Number – Single Precision
- c. BubblerDrip – Irrigation system is composed of bubbler and drip systems
  - i. Number – Integer (1 = Yes, 0 = No)

- d. BubblerDripSpray – Irrigation system is composed of bubbler, drip, and spray systems
  - i. Number – Integer (1 = Yes, 0 = No)
- e. Bubblers – Irrigation system is composed of bubblers
  - i. Number – Integer (1 = Yes, 0 = No)
- f. BubblerSpray – Irrigation system is composed of bubbler and spray systems
  - i. Number – Integer (1 = Yes, 0 = No)
- g. Drip – Irrigation system is composed of drip systems
  - i. Number – Integer (1 = Yes, 0 = No)
- h. DripOff – Irrigation system is composed of drip systems with one or more other irrigation zones turned off
  - i. Number – Integer (1 = Yes, 0 = No)
- i. DripMicro – Irrigation system is composed of drip and micro-spray systems
  - i. Number – Integer (1 = Yes, 0 = No)
- j. DripPopup – Irrigation system is composed of drip and popup spray systems
  - i. Number – Integer (1 = Yes, 0 = No)
- k. DripSpray – Irrigation system is composed of drip and spray systems
  - i. Number – Integer (1 = Yes, 0 = No)
- l. Hose – Irrigation is done with a hose
  - i. Number – Integer (1 = Yes, 0 = No)
- m. Microspray – Irrigation system is composed of micro-spray systems
  - i. Number – Integer (1 = Yes, 0 = No)
- n. Sprays – Irrigation system is composed of spray systems
  - i. Number – Integer (1 = Yes, 0 = No)
- o. BubblerDripPopup – Irrigation system is composed of bubbler, drip, and popup spray systems
  - i. Number – Integer (1 = Yes, 0 = No)
- p. DripMicroPopup – Irrigation system is composed of drip micro-spray and popup spray systems
  - i. Number – Integer (1 = Yes, 0 = No)
- q. DripPopupSpray – Irrigation system is composed of drip, popup spray, and spray systems
  - i. Number – Integer (1 = Yes, 0 = No)
- r. DripPopupRotor – Irrigation system is composed of drip, popup spray, and rotor systems
  - i. Number – Integer (1 = Yes, 0 = No)
- s. DripLaser – Irrigation system is composed of drip and laser tube systems
  - i. Number – Integer (1 = Yes, 0 = No)
- t. DripSoaker – Irrigation system is composed of drip and soaker hose systems
  - i. Number – Integer (1 = Yes, 0 = No)
- u. DripTurfBubbler – Irrigation system is composed of drip and turf bubbler systems
  - i. Number – Integer (1 = Yes, 0 = No)
- v. DripFountain – Irrigation system is composed of drip systems, and a fountain refill is controlled with the irrigation clock
  - i. Number – Integer (1 = Yes, 0 = No)

**13. tblMulches – 715 Records, mulch and weed barrier information**

- a. ClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. txtMulch – Typical type of mulch
  - i. Text - 18
- c. txtMulchSize – Typical size of mulch
  - i. Text - 50
- d. txtMulchColor – Typical color of mulch
  - i. Text - 6
- e. nstMulchDepth – Depth of mulch in inches
  - i. Number – Single Precision
- f. yntWeeds – Indicates if excessive weeds are present
  - i. Boolean
- g. yntSlope – Is a steep slope present?
  - i. Boolean
- h. yntTraffic – Is there heavy traffic in landscape?
  - i. Boolean
- i. yntAlkali – Indicates if excessive alkali deposits present at surface.
  - i. Boolean
- j. txtBarrierType – Type of weed barrier
  - i. Text – 20
- k. txtBarrierColor – Color of weed barrier
  - i. Text – 6
- l. yntBarrierShowing – Is the barrier showing at surface?
  - i. Boolean
- m. txtWear – Extent of wear
  - i. Text – 6
- n. txtLocationType – Wear location type
  - i. Text – 16

**14. tblMainmeterConsumption – 4318 Records, Gallons used per customer per month as measured by the property's main service meter.**

- a. nltClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. nitYear
  - i. Number – Integer
  - ii. Primary Key
- c. txtEntity – Indicates which water provider services the customer
  - i. Text – 5
- d. nstJan – January consumption in gallons
  - i. Number – Single Precision
- e. nstFeb – February consumption in gallons
  - i. Number – Single Precision



- f. nstMar – March consumption in gallons
  - i. Number – Single Precision
- g. nstApr – April consumption in gallons
  - i. Number – Single Precision
- h. nstMay – May consumption in gallons
  - i. Number – Single Precision
- i. nstJun – June consumption in gallons
  - i. Number – Single Precision
- j. nstJul – July consumption in gallons
  - i. Number – Single Precision
- k. nstAug – August consumption in gallons
  - i. Number – Single Precision
- l. nstSep – September consumption in gallons
  - i. Number – Single Precision
- m. nstOct – October consumption in gallons
  - i. Number – Single Precision
- n. nstNov – November consumption in gallons
  - i. Number – Single Precision
- o. nstDec – December consumption in gallons
  - i. Number – Single Precision
- p. nstTotal – Total annual consumption in gallons
  - i. Number – Single Precision

**15. tbl2001PropAreasOK4 – 673 Records,** Property area information as recorded for the year 2001.

- a. nltClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. Converted – Area converted from turf to xeriscape. Refers to “XS” Participants only.
  - i. Number – Single Precision
- c. Pool – Pool area if applicable
  - i. Number – Single Precision
- d. GardenMon – Garden area where irrigation is being monitored by the submeter
  - i. Number – Single Precision
- e. GardenUnmon – Garden area where irrigation is unmonitored by the submeter
  - i. Number – Single Precision
- f. Other – Square footage of other undeveloped property area. No irrigation, plants or hardscape present.
  - i. Number – Single Precision
- g. Study – Total xeriscape area where irrigation is monitored by the submeter. Applies to XS participant only.
  - i. Number – Single Precision
- h. TurfMon – Square footage of turf grass where irrigation is monitored by the submeter.
  - i. Number – Single Precision

- i. TurfUnmon – Square footage of turf area where the irrigation is not monitored by the submeter
    - i. Number – Single Precision
  - j. XeriMon – Square footage of xeriscape where irrigation is monitored by the submeter. (Applies to xeriscape study Group)
    - i. Number – Single Precision
  - k. XeriUnmon – Square footage of xeriscape area where the irrigation is not monitored by the submeter.
    - i. Number – Single Precision
  - l. TotalEvaporative – Total of all landscape areas plus pool area.
    - i. Number – Single Precision
  - m. TotalLandscape – Total of all landscape areas.
    - i. Number – Single Precision
  - n. dt2001SR – Date of 2001 follow-up site visit
    - i. Date / Time
  - o. AgeOfXeriscape – Age of xeriscape in days calculated by the difference between the post submeter installation inspection and the final 2001 follow-up site visit.
    - i. Number – Long Integer
  - p. TotalXeriArea – Total of all xeriscaped areas
    - i. Number – Single Precision
  - q. TotalGarden – Total of all garden areas
    - i. Number – Single Precision
  - r. TotalTurf – Total of all Turf areas
    - i. Number – Single Precision
  - s. PctGarden – Percent of total landscape area in garden
    - i. Number – Single Precision
  - t. PctXeri – Percent of total landscape in xeriscape
    - i. Number – Single Precision
  - u. PctTurf – Percent of total landscape area in turf
    - i. Number – Single Precision
  - v. PctOther – Percent of total landscape in other non-landscaped area
    - i. Number – Single Precision
  - w. PctPool – Percent of total landscape in pool
    - i. Number – Single Precision
  - x. PctXeriRank – Xeriscape study participants were divided into ten percent ranges based upon percentage of landscape in xeriscape and given a ranking.
    - i. Number – Long Integer
16. **tblTurfOnlySubMonthly – 107 Records**, monthly submeter consumption data and per square foot usage for turf study group of participants. Note – this usage is limited to those TS participants where ONLY turf was irrigated with submeter-monitored usage.
- a. ClientID – SNWA Customer identification number
    - i. Number – Long Integer
    - ii. Primary Key
  - b. Year
    - i. Number – Integer

- c. Entity – Water purveyor that serves the customer
  - i. Text – 5
- d. FileQuality – Quality rating of file information
  - i. Text – 10
- e. Status – Customer status
  - i. Text – 7
- f. TurfMon – Square feet of grass where irrigation is monitored by the submeter
  - i. Number – Single
- g. JanCons – January submeter consumption in gallons
  - i. Number – Single
- h. FebCons – February submeter consumption in gallons
  - i. Number – Single
- i. MarCons – March submeter consumption in gallons
  - i. Number – Single
- j. AprCons – April submeter consumption in gallons
  - i. Number – Single
- k. MayCons – May submeter consumption in gallons
  - i. Number – Single
- l. JunCons – June submeter consumption in gallons
  - i. Number – Single
- m. JulCons – July submeter consumption in gallons
  - i. Number – Single
- n. AugCons – August submeter consumption in gallons
  - i. Number – Single
- o. SepCons – September submeter consumption in gallons
  - i. Number – Single
- p. OctCons – October submeter consumption in gallons
  - i. Number – Single
- q. NovCons – November submeter consumption in gallons
  - i. Number – Single
- r. DecCons – December submeter consumption in gallons
  - i. Number – Single
- s. JanGalSF – Gallons used per square foot of turf in January
  - i. Number – Single
- t. FebGalSF – Gallons used per square foot of turf in February
  - i. Number – Single
- u. MarGalSF – Gallons used per square foot of turf in March
  - i. Number – Single
- v. AprGalSF – Gallons used per square foot of turf in April
  - i. Number – Single
- w. MayGalSF – Gallons used per square foot of turf in May
  - i. Number – Single
- x. JunGalSF – Gallons used per square foot of turf in June
  - i. Number – Single
- y. JulGalSF – Gallons used per square foot of turf in July
  - i. Number – Single

- z. AugGalSF – Gallons used per square foot of turf in August
  - i. Number – Single
- aa. SepGalSF – Gallons used per square foot of turf in September
  - i. Number – Single
- bb. OctGalSF – Gallons used per square foot of turf in October
  - i. Number – Single
- cc. NovGalSF – Gallons used per square foot of turf in November
  - i. Number – Single
- dd. DecGalSF – Gallons used per square foot of turf in December
  - i. Number – Single

**17. tblTurfOnlySubYearly – 107 Records**, yearly submeter consumption data and per square foot usage for turf study group of participants. Note – this usage is limited to those TS participants where ONLY turf was irrigated with submeter-monitored usage.

- a. ClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. Year
  - i. Number – Integer
  - ii. Primary Key
- c. Entity – Water purveyor that serves the customer
  - i. Text – 5
- d. TurfMon – Square feet of grass where irrigation is monitored by the submeter
  - i. Number – Single
- e. GalSqFt – Gallons used per square foot of turf per year
  - i. Number – Single
- f. YearlyCons – Total submetered consumption for the year.
  - i. Number – Single
- g. FileQuality – Quality rating of file information
  - i. Text - 8
- h. Status – Customer status
  - i. Text – 7

**18. tblXeriOnlySubMonthly – 1550 Records**, monthly submeter consumption data and per square foot usage for xeriscape study group of participants. Note – this usage is limited to those XS participants where ONLY xeriscape was irrigated with submeter-monitored usage.

- a. ClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. Year
  - i. Number – Integer
  - ii. Primary Key
- c. Entity – Water purveyor that serves the customer
  - i. Text – 5

- d. ConvNew – Indicates if the property’s xeriscape was a new installation or a conversion of grass to xeriscape.
  - i. Text – 4
- e. Status – Customer status
  - i. Text – 7
- f. FileQuality – Quality rating of file information
  - i. Text – 10
- g. XeriMon – Square feet of xeriscape where irrigation is monitored by the submeter
  - i. Number – Single Precision
- h. JanCons – January submeter consumption in gallons
  - i. Number – Single Precision
- i. FebCons – February submeter consumption in gallons
  - i. Number – Single Precision
- j. MarCons – March submeter consumption in gallons
  - i. Number – Single Precision
- k. AprCons – April submeter consumption in gallons
  - i. Number – Single Precision
- l. MayCons – May submeter consumption in gallons
  - i. Number – Single Precision
- m. JunCons – June submeter consumption in gallons
  - i. Number – Single Precision
- n. SepCons – September submeter consumption in gallons
  - i. Number – Single Precision
- o. OctCons – October submeter consumption in gallons
  - i. Number – Single Precision
- p. NovCons – November submeter consumption in gallons
  - i. Number – Single Precision
- q. DecCons – December submeter consumption in gallons
  - i. Number – Single Precision
- r. JanGalSF – Gallons used per square foot of xeriscape in January
  - i. Number – Single
- s. FebGalSF – Gallons used per square foot of xeriscape in February
  - i. Number – Single
- t. MarGalSF – Gallons used per square foot of xeriscape in March
  - i. Number – Single
- u. AprGalSF – Gallons used per square foot of xeriscape in April
  - i. Number – Single
- v. MayGalSF – Gallons used per square foot of xeriscape in May
  - i. Number – Single
- w. JunGalSF – Gallons used per square foot of xeriscape in June
  - i. Number – Single
- x. JulGalSF – Gallons used per square foot of xeriscape in July
  - i. Number – Single
- y. AugGalSF – Gallons used per square foot of xeriscape in August
  - i. Number – Single

- z. SepGalSF – Gallons used per square foot of xeriscape in September
  - i. Number – Single
- aa. OctGalSF – Gallons used per square foot of xeriscape in October
  - i. Number – Single
- bb. NovGalSF – Gallons used per square foot of xeriscape in November
  - i. Number – Single
- cc. DecGalSF – Gallons used per square foot of xeriscape in December
  - i. Number – Single

**19. tblXeriOnlySubYearly – 1550 Records**, yearly submeter consumption data and per square foot usage for xeriscape study group of participants. Note – this usage is limited to those XS participants where ONLY xeriscape was irrigated with submeter-monitored usage.

- a. ClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. Year
  - i. Number – Integer
  - ii. Primary Key
- c. Entity – Water purveyor that serves the customer
  - i. Text – 5
- d. ConvNew – Indicates if the property’s xeriscape was a new installation or a conversion of grass to xeriscape.
  - i. Text – 4
- e. XeriMon – Square feet of xeriscape where irrigation is monitored by the submeter
  - i. Number – Single Precision
- f. YearlyCons– Total submetered consumption for the year.
  - i. Number – Single
- g. GalSqFt – Gallons used per square foot of monitored xeriscape per year
  - i. Number – Single
- h. FileQuality – Quality rating of file information
  - i. Text – 10
- i. Status – Customer status
  - i. Text – 7

**20. tblPlantList – 538 Records**, list of plants used to verify xeriscape participant’s compliance with minimum canopy standards for program participation and classification of landscape plants in subsequent follow-up visits.

- a. PlantID
  - i. Number – Long Integer
  - ii. Primary Key
- b. Genus
  - i. Text - 50
- c. Species
  - i. Text - 50

- d. Var/Cult – Variety or cultivar of plant
  - i. Text - 50
- e. Common Name
  - i. Text - 50
- f. Width – Expected mature width of the plant
  - i. Number - Single
- g. Height – Expected mature height of the plant
  - i. Number - Integer
- h. Plant Habit – Type of plant (shrub, tree, etc.)
  - i. Text - 50
- i. H2OUse – Rated plant water needs.
  - i. Text – 50

**21. tbl2001HomeSales – 45 Records**, data provided by SalesTraq. Information related to home sales in Southern Nevada area in the year 2001 by zip code.

- a. Zipcode
  - i. Text – 5
  - ii. Primary Key
- b. NumberSold – Number of homes sold in zip code
  - i. Number – Single Precision
- c. MedianPrice – Median price of homes sold in zip code
  - i. Number – Single Precision
- d. AvgPrice – Average price of homes sold in zip code.
  - i. Number – Single Precision
- e. AvgPricePerSqFt – Average Price per square foot of homes sold in zip code.
  - i. Number – Single Precision
- f. AvgSize – Average size of homes sold in zip code.
  - i. Number – Single Precision
- g. Volume – Total value of homes sold in zip code
  - i. Number – Single Precision
- h. AvgAge – Average age of homes sold in zip code
  - i. Number – Single Precision

**22. tblMeterInfo – 716 Records**

- a. ClientID – SNWA Customer identification number
  - i. Number – Long Integer
  - ii. Primary Key
- b. MeterNum – Serial number stamped on submeter by manufacturer
  - i. Text – 50
- c. Installed – Date submeter was installed by contractor
  - i. Date/Time
- d. Cost – Cost of meter installation
  - i. Number – Single Precision
- e. RetrofitNum – AS/400 account number assigned to submeter
  - i. Number – Long Integer

- f. Location – approximate location of submeter on site.
  - i. Memo



#### APPENDIX 4: INFORMATION ON SINGLE-FAMILY RESIDENTIAL WATER BILL MODEL

A model was used to explore the differences in water consumption charges for a typical fifth decile in consumption LUC 110 property (single-family home) and one doing an average-size conversion. The model assumes the properties are in the Las Vegas Valley Water District’s service area and subject to its regular service rules. A typical 5/8-inch-meter size was assumed (meter size in large part determines rate per consumption unit). Rates for each tier and the size of the tier rate block appear below in the screen shot of the actual modeling processes for the model used in this report. As demonstrated, within a given billing cycle the rate for the first 5,000 gallons is \$1.05/kgal, the *next* 5,000 gallons after the initial 5000 costs \$1.75/kgal, the next 10,000 gallons after these first 10,000 gallons is \$2.38/kgal and so on (for billing purposes, the utility rounds to the nearest thousand gallons). In addition to the direct charges for the water, SNWA purveyor members bills commonly include a service charge, a commodity charge, and a reliability charge and these are reflected in the model below, so that the outputs are reflective of actual bills. A 30-day billing cycle was assumed.

In practical terms, the calculation of outputs in the model and the savings is derived by multiplying the expected average savings per square foot per month that would be yielded by a conversion (as calculated from Table 18) by the average-size conversion and then subtracting this from the fifth-decile consumption level. This yielded the costs with having done the conversion (below called “Total Bill”). In contrast, the cost without doing the conversion (i.e., “Average Fifth-Decile bill without reduction”) is shown under the “did conversion” scenario. The difference between these, highlighted in red, is the anticipated monthly bill savings yielded from having completed the conversion project.

#### Water Bill Calculator Screen Shot

Meter Size		Residential		Percent Reduction		0%		Select Entity		LVVWD Current		Average SF WSL Conversion (n=3651)		
5/8												1615.0		
		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual Totals
		10670	9369	12255	16174	21107	23447	25084	24519	21789	18112	13738	11795	208,057
		1.81	2.09	2.45	4.64	7.74	8.78	9.62	7.96	6.25	4.54	3.02	1.56	5th Decile average consumption
		2925	3377	3959	7497	12506	14187	15544	12862	10099	7336	4880	2521	97691
		Average Gallons Per SqFt Reduction												
		Avg Gal / SqFt Reduction * Avg SF WSL Conversion												
		Avg 5th Decile Cons - (Reduction * Avg WSL Conversion)												
Tier Rates	Tier Blocks	8	6	8	9	9	9	10	12	12	11	9	9	110366
1.05	5	5.25	5.25	5.25	5.25	5.25	5.25	5.25	5.25	5.25	5.25	5.25	5.25	63.00
1.75	5	4.80	1.74	5.77	6.43	6.30	7.46	7.94	8.75	8.75	8.75	6.75	7.48	80.93
2.38	10								3.94	4.02	1.85			9.81
3.02	20													0.00
Water Cost		10.05	6.99	11.02	11.68	11.55	12.71	13.19	17.94	18.02	15.85	12.00	12.73	153.74
Service														
3.669 Charge		3.67	3.67	3.67	3.67	3.67	3.67	3.67	3.67	3.67	3.67	3.67	3.67	44.03
Commodity														
0.05 Charge		0.39	0.30	0.41	0.43	0.43	0.46	0.48	0.58	0.58	0.54	0.44	0.46	5.52
Reliability														
0.25% Surcharge		0.04	0.03	0.04	0.04	0.04	0.04	0.04	0.06	0.06	0.05	0.04	0.04	0.51
Total Bill		14.15	10.98	15.14	15.83	15.69	16.88	17.38	22.25	22.33	20.11	16.16	16.90	203.79
		19.85	17.08	23.71	33.25	45.98	53.18	58.22	56.48	48.08	37.98	27.32	22.59	443.72
		5.70	6.09	8.57	17.42	30.29	36.30	40.84	34.23	26.25	17.87	11.17	5.69	239.93
		Average 5th Decile bill without reduction												
		Total Savings with Red												

## **APPENDIX 5: INFORMATION ON HOMEOWNER PERSPECTIVE MODEL**

The model is a dynamic Net Present Value Model that calculates the NPV of the project in future years. It does this by computing the difference in the yield by converting to xeriscape to the costs (water and maintenance) incurred by keeping turfgrass over the years.

“Conversion cost” and “awarded incentive” are products of the associated rates and the square feet converted. These are onetime costs. The “interest rate” is the discount or alternative rate (i.e., the rate associated with the loss incurred by spending money on the conversion rather than placing it in an interest-bearing account). The “average yearly rate increase” is the long-time average increase in water costs. “Yearly maintenance savings” is a product of the “Labor Savings” and “Direct Maintenance” variables (which are themselves calculated in a manner similar to “awarded incentive,” however, these savings are yielded each year). “Average total bill savings for a year” is not automatically calculated, but entered either by use of real data or modeled bill savings (see Appendix 4). Model Outputs are “NPV” and “Year.” Year 0 is the year of the conversion.

This model can directly yield the payback time with and without the incentive. By iterative process one can then develop what the input variables values would need to provide for a positive NPV at a given year. This is how the values for the third and fifth-year ROIs were developed for Figure 15. Example inputs and outputs are given below. In this case, at \$1.00 per square foot, the conversion reached a positive NPV between years one and two.

In terms of yielding the actual data in Table 15, the following were used as data sources:

“Square Feet Converted” – This was the average conversion size for SNWA’s Water Smart Landscapes Program in early 2004.

“Incentive Level” – This was the \$1.00 per square foot incentive level for almost all single-family conversion projects in SNWA’s Water Smart Landscapes Program in early 2004 (also see Appendix 5).

“Conversion cost” – This was the conversion cost as revealed by survey. This was one of the variables that were modified to reflect whether or not one did the conversion themselves or utilized contract assistance. Rates for each of these scenarios were developed based on compilation of receipts from both types of installations.

“average total bill savings for a year” – This was the yearly savings as provided by a model of the Las Vegas Valley Water District for a LUC 110 property in the fifth decile (mid-range) of consumption (see Appendix 4 for details on this model).

“interest rate” – This was the interest rate of a home equity loan in early February 2004.

“average yearly rate increase” – This is the average yearly rate increase for the Las Vegas Valley Water District over the long term. In practice, the District has often gone significant periods of

time without a rate increase and then increased them much more than 3%, but this was the most practical method of doing the calculation for purposes of creating the model.

“Labor Savings” – This was adapted from Hessling<sup>12</sup> (2001). This savings was effectively turned on or off to see the impacts of the situations when labor savings are and are not realized. See text for additional information.


“Direct Maintenance” – This rate was derived from the maintenance survey data and is per Hessling<sup>12</sup> (2001).

Examples of Homeowner Perspective Model Inputs and Outputs

<b>Inputs:</b>	Type	NPV	Year
Square Feet Converted	1616		
Incentive level	\$1.00		
Conversion cost:	\$1.37		
conversion cost:	\$2,213.92	<b>(\$2,070.88)</b>	0
average total bill savings for a year:	\$240.00	<b>(\$636.58)</b>	1
awarded incentive:	\$1,616.00	<b>\$751.63</b>	2
interest rate:	6.32%	<b>\$2,095.24</b>	3
average yearly rate increase	3.00%	<b>\$3,395.67</b>	4
		<b>\$4,654.31</b>	5
Labor Savings	\$0.20		
Labor Savings	\$323.20		
Direct Maintenance	\$0.11		
Direct Maintenance	\$177.76		
Yearly maintenance savings	\$500.96		

## APPENDIX 6: INFORMATION ON SNWA'S WATER SMART LANDSCAPES PROGRAM

Growth of Program:



See Program Application (following)

